

Standard Errors When a Regressor is Randomly Assigned*

Denis Chetverikov[†]
UCLA

Jinyong Hahn[‡]
UCLA

Zhipeng Liao[§]
UCLA

Andres Santos[¶]
UCLA

March 7, 2026

Abstract

We show that the OLS variance formula simplifies when values of the regressor of interest are assigned randomly and independently of other regressors. When the regressor of interest is independent not only of other regressors but also of the error term, the homoskedastic variance formula is valid even if regression errors are correlated. When the error term is heteroscedastic with respect to the regressor of interest, the variance formula has to be adjusted not only for heteroscedasticity but also for correlation structure of the error term. However, even in the latter case, some simplifications are possible.

JEL Classification: C14, C31, C32

Keywords: Cluster Robust Inference; Randomized Control Trial; Bartik Regression

1 Introduction

Textbook discussion of linear regression usually begins with a standard model of the form $\mathbf{Y} = \mathbf{X}\theta + \epsilon$, where it is assumed that $\mathbf{X}$ is a nonstochastic matrix (with full column rank) of regressors and the error vector ϵ has mean zero and variance matrix proportional to an identity matrix. As is well known, such an assumption justifies the formula $s^2 (\mathbf{X}^\top \mathbf{X})^{-1}$ as an estimator of the variance of the OLS estimator, where s^2 is equal to the sum of squares of the estimated residuals divided either by the sample size or the degrees of freedom. This formula is easy to use but, as is typically taught, may not be valid if the variance matrix Ω of the error vector ϵ is not proportional to an identity matrix. In such cases, the variance of the OLS estimator should be based on the formula $(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \Omega \mathbf{X} (\mathbf{X}^\top \mathbf{X})^{-1}$ to reflect the heteroscedasticity and dependence structure of the error

*We thank Steve Pischke for helpful comments.

[†]Department of Economics, UCLA, Los Angeles, CA 90095-1477 USA. Email: chetverikov@econ.ucla.edu

[‡]Department of Economics, UCLA, Los Angeles, CA 90095-1477 USA. Email: hahn@econ.ucla.edu

[§]Department of Economics, UCLA, Los Angeles, CA 90095-1477 USA. Email: zhipeng.liao@econ.ucla.edu

[¶]Department of Economics, UCLA, Los Angeles, CA 90095-1477 USA. Email: andres@econ.ucla.edu

vector. An important practical challenge in implementing such an approach is that the matrix Ω may be hard to estimate if the dependence structure of the error vector ϵ is unknown. In this paper, we study the implications for the variance of OLS estimators of having a regressor of interest whose values are i.i.d. across units/time periods and are independent of values of other regressors. Our motivating applications include certain completely randomized controlled trials and related quasi-experimental designs such as shift-share regressions. The main finding of the paper is that variance estimation in this case is often simplified, sometimes substantially.

Let $\mathbf{D}$ be the column of $\mathbf{X}$ corresponding to the regressor of interest and let $\mathbf{W}$ be the remaining columns of $\mathbf{X}$; i.e. columns corresponding to controls. Our first main result shows that when the vector $\mathbf{D}$ has i.i.d. components and is *strongly exogenous* in the sense of being independent not only of $\mathbf{W}$ but also of ϵ , the OLS estimator is asymptotically normally distributed and the formula $s^2 (\mathbf{X}^\top \mathbf{X})^{-1}$ actually yields a valid variance estimator for the coefficient of $\mathbf{D}$ even if Ω is not proportional to the identity matrix. This result, which superficially contradicts the lessons of elementary linear regression analysis, is due to the randomness of the $\mathbf{X}$ matrix in our analysis. While the textbook analysis assumes away the randomness by conditioning on $\mathbf{X}$, we instead obtain our result by recognizing that the randomness of the $\mathbf{X}$ matrix delivers a suitable martingale structure.¹ We recognize that a version of this result in some simple contexts is well understood in the profession in the sense that many can anticipate such a result when the entire matrix of regressors is strongly exogenous; see references below. We go one step further, however, and establish our result for the case when (i) only *one* regressor is strongly exogenous (e.g., treatment in a randomized controlled trial); and/or (ii) the error vector is subject to some generalized dependence more complicated than what is commonly understood to be the cluster structure, e.g. temporal/spatial autocorrelation or a network structure. This result is important because it facilitates inference on the coefficient of the regressor of interest even if the researcher does not know the dependence structure of the error vector ϵ , which is useful from the pragmatic point of view. We emphasize, however, that while our conclusions hold for the coefficient corresponding to a strongly exogenous regressor, they need not hold for other coefficients in the regression.

Our second main result shows that when the vector $\mathbf{D}$ has i.i.d. components and is independent of $\mathbf{W}$ but ϵ is *conditionally heteroscedastic* with respect to $\mathbf{D}$, the formula $s^2 (\mathbf{X}^\top \mathbf{X})^{-1}$ is actually not valid and has to be adjusted not only for heteroscedasticity *but also*, rather surprisingly, for the dependence structure of the vector ϵ . Nevertheless, a simplified variance formula can often be used in this case as well. For example, conditional heteroscedasticity arises when the regression model $\mathbf{Y} = \mathbf{X}\theta + \epsilon$ is taken from the potential outcomes framework with heterogeneous treatment effects. If the researcher is concerned about clustering in this case, our results imply that it is

¹The validity of the formula $s^2 (\mathbf{X}^\top \mathbf{X})^{-1}$ does not mean that the variance estimators based on the formula $(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \Omega \mathbf{X} (\mathbf{X}^\top \mathbf{X})^{-1}$ are invalid; see Lemma 1 in the Appendix for the asymptotic equivalence of these estimators in our setting.

sufficient to adjust the variance formula for clustering of the treatment effects only.² In contrast, for example, there is no need to adjust the variance formula for clustering of the potential outcomes in any given treatment arm. We also note that neither of our results restrict or exclude conditional heteroscedasticity of ϵ with respect to $\mathbf{W}$.

In addition, we extend both results to the case when the values of the regressor of interest are independent only *across groups* of units/time periods, such as is the case in randomized controlled trials in which treatment assignment is determined at a group (e.g., school/village) level. We show that in the strongly exogenous case, it suffices to take into account only the within-group correlation of the error vector ϵ . In other words, it suffices to use variance estimators that are clustered at the level at which treatment is assigned. In the conditional heteroscedasticity case, the variance formula still requires some adjustments for both heteroscedasticity and dependence but often allows for some simplifications relative to the standard textbook formula mentioned above.

Our first main result and its extension to the group-level assignment are related to but different from those in Barrios, Diamond, Imbens, and Kolesar (2012), who came to the same conclusions for regressions without controls and in which a fixed fraction of units/clusters is randomly assigned to be treated. In a related but independent study, Zhang, Ding, Zhou, and Wang (2025) demonstrate a similar robustness of the classical standard error formula $s^2 (\mathbf{X}^\top \mathbf{X})^{-1}$, assuming that the rows of $\mathbf{X}$ are independent and independent of ϵ . Our setting differs in that we focus on inference for the coefficient on the key regressor $\mathbf{D}$, and we allow for general dependence in both the error term ϵ and the controls $\mathbf{W}$. To the best of our knowledge, however, there are no results in the literature related to our second main result.

Our analysis is also related to Abadie, Athey, Imbens, and Wooldridge (2023), who presented a new clustering framework that emphasizes a finite population perspective as well as interactions between the sampling and assignment parts of the data-generating process. They established in particular that there is no need to cluster when estimating the variance if the randomization is conducted at the individual level and there is no heterogeneity in the treatment effects. Our first main result echoes and complements their findings in the following aspects. First, unlike them, we do not impose a particular structure on the sampling process, which allows us to cover general forms of time series or even network dependence in addition to the cluster-type dependence. Second, our analysis goes beyond the binary treatment framework and accommodates a general strongly exogenous regressor as well as the inclusion of additional controls in the regression. In particular, we allow for general dependence structures in the control variables, which makes it ex-ante unclear at what level one should cluster. Third, we rely on a traditional asymptotic framework, which may make our analysis more familiar to the reader. We recognize, however, that the third aspect may be

²When the regression model $\mathbf{Y} = \mathbf{X}\theta + \epsilon$ is taken from the potential outcomes framework with heterogeneous treatment effects, the case of strongly exogenous regressor corresponds to the assumption of constant treatment effects.

a weakness in the sense that our framework is unable to address the finite population adjustment that plays an important role in Abadie, Athey, Imbens, and Wooldridge (2023). Finally, we note Bloom (2005) who considered a random effects type cluster structure and produced a variance estimator for the simple difference of means estimator that is quoted in Duflo, Glennerster, and Kremer (2007). His equation (4.3), which is presented without proof, is in fact a special case of the $s^2 (\mathbf{X}^\top \mathbf{X})^{-1}$ formula. The cluster structure that he analyzed has a built-in correlation among observations, and as such, it would be tempting to think that variance estimation would require Moulton (1986)’s clustering formula – a conclusion that can be motivated if inference is to be conditioned on the $\mathbf{X}$ matrix. Hence, in our view, his equation (4.3) can only be motivated by explicitly recognizing the randomness of the $\mathbf{X}$ matrix.

Our results are not particularly difficult to derive. On the other hand, we are unaware of any systematic discussion of results along this line in the literature besides Barrios, Diamond, Imbens, and Kolesar (2012), Abadie, Athey, Imbens, and Wooldridge (2023) and Zhang, Ding, Zhou, and Wang (2025), especially in models where the control variables and treatment variables are both present. Our results have convenient pragmatic implications, which we hope are helpful to some empirical researchers.

Finally, we establish a connection between our analysis and a recent literature on shift-share regressions that attributes identification to the random assignment of shocks (Adao, Kolesár, and Morales, 2019; Borusyak, Hull, and Jaravel, 2022). In particular, our analysis suggests a novel two step estimator for these models. In the first step, we run OLS regressions of both the outcome of interest and the explanatory variables on the shares. In the second step, we estimate the shift-share regression coefficients by running an OLS regression of the first stage estimated coefficients for the outcome variable on the estimated first stage coefficients for the explanatory variable. The motivation for this procedure is that the regressor of interest in the second stage regression consists of the shocks in the shift-share design, which are often assumed to be i.i.d. and independent of other variables. As a result, the second stage regression has the structure that is the focus of our analysis and our results deliver a simple formula for the asymptotic variance of the proposed estimator.

Outline. We present the basic intuition underlying our results in Section 2. The intuitive discussion there suggests that asymptotic normality and the formula $s^2 (\mathbf{X}^\top \mathbf{X})^{-1}$ are valid under fairly general dependence structure in ϵ provided that the randomness of $\mathbf{X}$ generates a suitable martingale structure in $\mathbf{X}^\top \epsilon$. We also explain how conditional heteroscedasticity breaks down this martingale structure. We formalize our discussion in Section 3, where our main restriction on the dependence of ϵ is that it be weak enough for its sample average to converge in probability to zero – a condition that further emphasizes that our analysis is driven by the randomness in $\mathbf{X}$ and not ϵ . We provide an extension to the case of group-level random assignment in Section 4. We discuss shift-share regressions in Section 6.

Notation. We use K to denote a generic strictly positive constant that may change from place to place but is independent of the sample size n . For any positive integer k , let $\mathbf{I}_k$, $\mathbf{1}_{k \times 1}$, and $\mathbf{0}_{k \times 1}$ denote the $k \times k$ identity matrix, $k \times 1$ vector of ones, and $k \times 1$ vector of zeros. For any real square matrix A , $\lambda_{\min}(A)$ and $\lambda_{\max}(A)$ denote its smallest and largest eigenvalues. We use $A \equiv B$ to denote that A is defined as B , and $(A_j)_{j \leq J}$ to denote the matrix composed by sequentially stacking matrices $A_1, \dots, A_J$ with equal number of columns.

2 Intuition

In this section, we provide intuition for the validity of the formula $s^2 (\mathbf{X}^\top \mathbf{X})^{-1}$ in the case of strongly exogenous regressors. We first consider the case of a simple univariate regression model and then extend the result to the case of a multivariate regression model. At the end of this section, we explain complications arising from conditional heteroscedasticity and how they break the validity of the formula $s^2 (\mathbf{X}^\top \mathbf{X})^{-1}$.

2.1 Case of Strong Exogeneity

We start by considering a simple univariate linear regression time series model in which we have

$$y_i = d_i \beta + \varepsilon_i, \quad i = 1, \dots, n, \quad (1)$$

where d_i is assumed to have zero mean for simplicity in this subsection, and $(\varepsilon_i)_{i \leq n}$ is a second-order, possibly autocorrelated, stationary time series – we employ the index i , rather than t , to emphasize that our analysis is not confined to the time series context. We depart from the textbook time series model by assuming that the regressors $(d_i)_{i \leq n}$ are: (i) Independent and identically distributed (i.i.d.), and (ii) Strongly exogenous in the sense that $(d_i)_{i \leq n}$ is independent of the time series process $(\varepsilon_i)_{i \leq n}$.

As is well-known, the least squares estimator $\hat{\beta}$ of β in model (1) satisfies

$$\sqrt{n}(\hat{\beta} - \beta) = \frac{n^{-1/2} \sum_{i=1}^n d_i \varepsilon_i}{n^{-1} \sum_{i=1}^n d_i^2}. \quad (2)$$

In many standard time series textbooks, the asymptotic distribution of $\hat{\beta}$ is thus derived by imposing sufficiently strong conditions to ensure that the score $n^{-1/2} \sum_{i=1}^n d_i \varepsilon_i$ is asymptotically normal and the Hessian $n^{-1} \sum_{i=1}^n d_i^2$ converges in probability to a non-stochastic matrix. In order to derive the standard error for $\hat{\beta}$, we therefore only need a consistent estimator of the long-run variance of the score; i.e., a heteroscedasticity and autocorrelation consistent (HAC) variance estimator, such as those proposed by Newey and West (1987) and Andrews (1991).

On the other hand, if the regressors $(d_i)_{i \leq n}$ are i.i.d. and strongly exogenous with mean zero, the independence of $(d_i)_{i \leq n}$ and $(\varepsilon_i)_{i \leq n}$ implies that for any $1 \leq i_1, i_2 \leq n$ with $i_1 \neq i_2$, we must

have

$$\mathbb{E}[d_{i_1}\varepsilon_{i_1}d_{i_2}\varepsilon_{i_2}] = \mathbb{E}[d_{i_1}]\mathbb{E}[d_{i_2}]\mathbb{E}[\varepsilon_{i_1}\varepsilon_{i_2}] = 0,$$

and also $\mathbb{E}[(d_i\varepsilon_i)^2] = \mathbb{E}[d_i^2]\mathbb{E}[\varepsilon_i^2]$ for all $1 \leq i \leq n$. Hence, as long as some version of the central limit theorem is applicable to the score $n^{-1/2} \sum_{i=1}^n d_i\varepsilon_i$ and a law of large numbers is applicable to the Hessian $n^{-1} \sum_{i=1}^n d_i^2$, we can conclude that the asymptotic distribution of $\hat{\beta}$ is given by

$$\sqrt{n}(\hat{\beta} - \beta) \rightarrow_d N\left(0, \frac{\mathbb{E}[\varepsilon_i^2]}{\mathbb{E}[d_i^2]}\right).$$

In particular, statistical inference on β can be conducted as if it were not a time series model, i.e. using the formula $s^2(\mathbf{X}^\top \mathbf{X})^{-1}$, where $\mathbf{X} \equiv (d_i)_{i \leq n}$ in this case.

The main takeaway of the preceding example is that the strong exogeneity and i.i.d. nature of the regressors $(d_i)_{i \leq n}$ imply that the sequence $(d_i\varepsilon_i)_{i \leq n}$ is homoscedastic *even if* the errors $(\varepsilon_i)_{i \leq n}$ are arbitrarily autocorrelated. Is this simplification confined to the time series model? Our preceding discussion suggests that this is not the case. Indeed, provided the regressors $(d_i)_{i \leq n}$ are i.i.d., mean zero, and strongly exogenous, the score $n^{-1/2} \sum_{i=1}^n d_i\varepsilon_i$ has a built-in martingale structure vis-à-vis the filtration $\mathcal{F}_i$ generated by $\{d_j, \varepsilon_j\}_{j \leq i}$ because:

$$\mathbb{E}[d_i\varepsilon_i | \mathcal{F}_{i-1}] = \mathbb{E}[\mathbb{E}[d_i | \mathcal{F}_{i-1}, \varepsilon_i] \varepsilon_i | \mathcal{F}_{i-1}] = \mathbb{E}[d_i]\mathbb{E}[\varepsilon_i | \mathcal{F}_{i-1}] = 0. \quad (3)$$

Therefore, assuming that the random pairs (d_i, ε_i) satisfy certain moment conditions, the martingale central limit theorem will be applicable regardless of the dependence structure of $(\varepsilon_i)_{i \leq n}$ and the long run variance of the score will reduce to $\mathbb{E}[d_i^2]\mathbb{E}[\varepsilon_i^2]$. In particular, the variance formula $s^2(\mathbf{X}^\top \mathbf{X})^{-1}$ will remain valid despite the dependence present in the variables $(\varepsilon_i)_{i \leq n}$. Thus, spatial correlation, network dependence, and/or a cluster structure in the variables $(\varepsilon_i)_{i \leq n}$ are all accommodated by the standard homoscedastic standard errors. Moreover, we note that a quick inspection at the preceding argument reveals that the assumptions we have imposed so far are stronger than necessary for the desired conclusion to hold.

We next build on our preceding discussion by considering the multivariate linear regression model

$$y_i = d_i\beta + w_i^\top \gamma + \varepsilon_i, \quad i = 1, \dots, n, \quad (4)$$

where d_i is a scalar regressor of interest, w_i is a d_w -vector of controls, and $\{(w_i^\top, \varepsilon_i)\}_{i \leq n}$ is a second-order, possibly autocorrelated, stationary time series satisfying $\mathbb{E}[w_i\varepsilon_i] = \mathbf{0}_{d_w \times 1}$ for all $1 \leq i \leq n$. We continue to assume that the regressors $(d_i)_{i \leq n}$ are i.i.d. with mean zero, and strongly exogenous in the sense that $(d_i)_{i \leq n}$ is independent of the time series process $(\varepsilon_i, w_i^\top)_{i \leq n}$. The parameter of interest continues to be β .

For this model, the Frisch-Waugh-Lovell theorem implies the least squares estimator $\hat{\beta}$ satisfies

$$\sqrt{n}(\hat{\beta} - \beta) = \frac{n^{-1/2} \sum_{i=1}^n (d_i - w_i^\top \hat{\alpha}) \varepsilon_i}{n^{-1} \sum_{i=1}^n (d_i - w_i^\top \hat{\alpha})^2}, \quad \hat{\alpha} \equiv \left(\sum_{i=1}^n w_i w_i^\top \right)^{-1} \sum_{i=1}^n w_i d_i. \quad (5)$$

Hence, under appropriate regularity conditions the estimator $\hat{\beta}$ admits the asymptotic expansion

$$\sqrt{n}(\hat{\beta} - \beta) = \frac{n^{-1/2} \sum_{i=1}^n (d_i - w_i^\top \alpha) \varepsilon_i}{n^{-1} \sum_{i=1}^n (d_i - w_i^\top \alpha)^2} + o_p(1), \quad \alpha \equiv (\mathbb{E}[w_i w_i^\top])^{-1} \mathbb{E}[w_i d_i]. \quad (6)$$

In particular, if $(d_i)_{i \leq n}$ is mean zero and independent of $(w_i^\top)_{i \leq n}$, then $\alpha = 0$ and the asymptotic expansion reduces to the univariate setting – i.e. the right-hand side of (6) is (asymptotically) equivalent to the right-hand side of (2). It therefore follows that the end result is the same as in the univariate case: The variance formula $s^2(\mathbf{X}^\top \mathbf{X})^{-1}$, where $\mathbf{X} \equiv (d_i, w_i^\top)_{i \leq n}$ in this case, remains valid for $\hat{\beta}$. We emphasize, however, that this formula is not necessarily justified for conducting inference on the coefficient γ .

As a preview of results in the next section, we again note that the conditions we have imposed so far are stronger than required. For instance, suppose that instead of demanding that the regressors $(d_i)_{i \leq n}$ and $(w_i^\top)_{i \leq n}$ be fully independent, we assume that they are related according to the model

$$d_i = w_i^\top \alpha + \eta_i,$$

for $(\eta_i)_{i \leq n}$ i.i.d. and independent of $(\varepsilon_i, w_i^\top)_{i \leq n}$. The asymptotic expansion in (6), combined with the same arguments employed in the univariate case, then continue to imply that

$$\sqrt{n}(\hat{\beta} - \beta) \rightarrow_d N\left(0, \frac{\mathbb{E}[\varepsilon_i^2]}{\mathbb{E}[\eta_i^2]}\right)$$

under mild moment conditions – i.e. when computing standard errors for $\hat{\beta}$ we can continue to pretend that the time series process fits the textbook homoscedastic model. Setting w_i to be a constant, for instance, reveals that the mean zero assumption on $(d_i)_{i \leq n}$ is superfluous.

2.2 Case of Conditional Heteroscedasticity

The preceding martingale argument relies crucially on two key assumptions: (i) The regressors $(d_i)_{i \leq n}$ are independent of each other, and (ii) The regressors $(d_i)_{i \leq n}$ are independent of the errors $(\varepsilon_i)_{i \leq n}$. A challenge to our martingale argument arises when $(d_i)_{i \leq n}$ and $(\varepsilon_i)_{i \leq n}$ are not independent. Within the potential outcome framework, for instance, this full independence requirement is violated in the presence of heterogenous treatment effects. More precisely, heterogenous treatment effects render ε_i conditionally heteroscedastic with respect to the treatment status d_i . Motivated by this observation, we also study a model in which $\varepsilon_i = \sum_{l=1}^L \sigma_l(d_i) \varepsilon_{l,i}^*$ with $e_i \equiv (\varepsilon_{l,i}^*)_{l \leq L}$ possibly correlated across i and l , but $(e_i)_{i \leq n}$ fully independent of $(d_i)_{i \leq n}$. In the potential outcome framework, with $d_i \in \{0, 1\}$ indicating treatment status, we would have

$$\varepsilon_i = (y_i(0) - \mathbb{E}[y_i(0)]) + d_i((y_i(1) - y_i(0)) - \mathbb{E}[y_i(1) - y_i(0)]), \quad (7)$$

where $y_i(d)$ denotes the potential outcome for unit i under treatment status $d \in \{0, 1\}$.³

To see the problem for the martingale structure in this model, observe that for any $1 \leq i_1, i_2 \leq n$, we now have

$$\begin{aligned}\mathbb{E}[d_{i_1}\varepsilon_{i_1}d_{i_2}\varepsilon_{i_2}] &= \mathbb{E}\left[d_{i_1}\sum_{l_1=1}^L\sigma_{l_1}(d_{i_1})\varepsilon_{l_1,i_1}^*d_{i_2}\sum_{l_2=1}^L\sigma_{l_2}(d_{i_2})\varepsilon_{l_2,i_2}^*\right] \\ &= \sum_{l_1=1}^L\sum_{l_2=1}^L\mathbb{E}[d_{i_1}\sigma_{l_1}(d_{i_1})]\mathbb{E}[d_{i_2}\sigma_{l_2}(d_{i_2})]\mathbb{E}[\varepsilon_{l_1,i_1}^*\varepsilon_{l_2,i_2}^*],\end{aligned}$$

which is not necessarily zero even if d_i 's are mean zero. The variance formula for the sum $\sum_{i \leq n} d_i \varepsilon_i$ therefore must include interactions terms as long as the random vectors $(\varepsilon_{l,i_1}^*)_{l \leq L}$ and $(\varepsilon_{l,i_2}^*)_{l \leq L}$ are correlated.

We will present a detailed analysis of the conditional heteroscedasticity case in Section 3 but the main takeaways from our results are: (i) The independence of $(d_i)_{i \leq n}$ from controls $(w_i^\top)_{i \leq n}$ still simplifies the asymptotic variance for $\hat{\beta}$; and (ii) Conditional heteroscedasticity yields a break in the martingale structure that requires us to adjust standard errors *not only* for heteroscedasticity, but *also* for correlation of the errors across units. For example, in the context of (7), our analysis implies the asymptotic variance of $\sqrt{n}(\hat{\beta} - \beta)$ equals (the probability limit of)

$$\begin{aligned}&\frac{1}{n\sigma_d^4}\sum_{i=1}^n\mathbb{E}[(d_i - \mathbb{E}[d_i])^2\varepsilon_i^2] \\ &+ \text{Var}\left(n^{-1/2}\sum_{i=1}^n(y_i(1) - y_i(0))\right) - n^{-1}\sum_{i=1}^n\text{Var}(y_i(1) - y_i(0)).\end{aligned}\quad (8)$$

In particular, we note that standard errors may need to be adjusted for correlation if we are concerned the treatment effects are correlated. On the other hand, the correlation between components of the vector $(y_i(0))_{i \leq n}$ plays no role for the standard errors, and neither does correlation between components of the vector $(y_i(1))_{i \leq n}$.

The validity or invalidity of the formula $s^2(\mathbf{X}^\top \mathbf{X})^{-1}$, which is discussed intuitively in this section, is formally established in the next section under suitable regularity conditions. Specifically, when the regressors $(d_i)_{i \leq n}$ are i.i.d. and independent of the errors $(\varepsilon_i)_{i \leq n}$, the formula $s^2(\mathbf{X}^\top \mathbf{X})^{-1}$ provides a valid standard error for $\hat{\beta}$. Combined with asymptotic normal approximation, this yields valid inference for β . The validity of $s^2(\mathbf{X}^\top \mathbf{X})^{-1}$ continues to hold in the presence of additional control variables, provided that the regressors $(d_i)_{i \leq n}$ are independent of these controls.

By contrast, when d_i is not independent of ε_i , as in the case of conditional heteroscedasticity, the formula $s^2(\mathbf{X}^\top \mathbf{X})^{-1}$ is no longer valid. In such settings, the underlying dependence structure

³The error term ε_i in (7) can be written as $\varepsilon_i = d_i(y_i(0) - \mathbb{E}[y_i(0)]) + (1 - d_i)(y_i(1) - \mathbb{E}[y_i(1)])$, which is a special case of our general specification $\varepsilon_i = \sum_{l=1}^L \sigma_l(d_i)\varepsilon_{l,i}^*$ with $L = 2$, $\varepsilon_{l,i}^* = y_i(l) - \mathbb{E}[y_i(l)]$ and $\sigma_l(d_i) = (-1)^l(d_i - l)$ for $l = 0, 1$.

of $(\varepsilon_i)_{i \leq n}$ must be explicitly accounted for in order to construct consistent standard errors and valid inference for $\hat{\beta}$.

These observations motivate the formal analysis that follows. In the next section, we introduce the precise assumptions governing the assignment mechanism and present the main theoretical results. Theorem 1, together with Corollary 1, rigorously establish the conditions under which the conventional variance formula is valid, while Theorem 2 characterizes the appropriate variance correction when it is not.

3 Main Theory

We now present a rigorous theory showing that the variance formula $s^2 (\mathbf{X}^\top \mathbf{X})^{-1}$ for the OLS estimator is valid in the strongly exogenous case even if the errors in the regression model are correlated. We also derive the variance formula in the case of conditional heteroscedasticity. In what follows, we suppose an outcome y_i satisfies

$$y_i = x_i^\top \theta + \varepsilon_i, \quad i = 1, \dots, n, \quad (9)$$

where $x_i = (d_i, w_i^\top)^\top$ is a vector of regressors, with d_i being a key regressor and w_i being a vector of controls (including a constant), $\theta = (\beta, \gamma^\top)^\top$ is a vector of parameters, with β being a parameter of interest and γ being a vector of nuisance parameters, and ε_i is an error term with mean zero. More explicitly, the regression model (9) can be rewritten as

$$y_i = d_i \beta + w_i^\top \gamma + \varepsilon_i, \quad i = 1, \dots, n. \quad (10)$$

We assume that the first component of the vector w_i is a (non-zero) constant, meaning that the regression model (10) contains the intercept term. For notational simplicity, we set $\mathbf{Y} \equiv (y_i)_{i \leq n}$, $\mathbf{D} \equiv (d_i)_{i \leq n}$, $\mathbf{W} \equiv (w_i^\top)_{i \leq n}$, $\mathbf{X} \equiv (x_i^\top)_{i \leq n}$, and $\boldsymbol{\epsilon} \equiv (\varepsilon_i)_{i \leq n}$.

Under a suitable exogeneity assumption on $(d_i, w_i^\top)^\top$, the unknown parameter $\theta \equiv (\beta, \gamma^\top)^\top$ can be estimated by OLS:

$$\hat{\theta} \equiv (\hat{\beta}, \hat{\gamma}^\top)^\top \equiv (\mathbf{X}^\top \mathbf{X})^{-1} (\mathbf{X}^\top \mathbf{Y}).$$

Standard estimators of the asymptotic variance of $\hat{\theta}$ rely on the asymptotic variance of the “score” $n^{-1/2} \sum_{i=1}^n (d_i, w_i^\top)^\top \varepsilon_i$, which may take a complicated form due to possible dependence between observations. This relationship makes standard error estimation and statistical inference challenging in practice. For instance, cluster-robust standard errors, such as those proposed by Moulton (1986), Liang and Zeger (1986), and Arellano (1987), are predicated on knowledge of the relevant group structure at which to cluster; see Hansen (2007), Ibragimov and Müller (2010) and Ibragimov and Müller (2016) for related discussion. Similarly, spatial standard errors, such as those proposed by Conley (1999), often require knowledge of a measure of “economic distance” that is

relates to the degree of dependence across observations. However, we next show that when $\mathbf{D}$ is independent of $\mathbf{W}$, such as in randomized control trials, the asymptotic variance for $\hat{\beta}$ simplifies significantly. As a result, estimation of standard errors and asymptotically valid inference simplify as well.

3.1 Case of Strong Exogeneity

We first study the case that $\mathbf{D}$ is independent of ϵ , and hence ϵ is conditionally homoscedastic with respect to $\mathbf{D}$. Together with independence of $\mathbf{D}$ from $\mathbf{W}$, this means that $\mathbf{D}$ is strongly exogenous. The conditional heteroscedasticity case is discussed later in this section. In the assumptions that follow, recall that K should be interpreted to denote a sufficiently large constant that can change from place to place but is independent of the sample size n .

Assumption 1. (i) The random variables d_i , $1 \leq i \leq n$, are i.i.d. with finite mean μ_d and variance σ_d^2 ; (ii) $(d_i)_{i \leq n}$ is independent of $(w_i^\top)_{i \leq n}$; (iii) there exists a constant $\delta_1 > 0$ such that $\max_{i \leq n} \mathbb{E}[|d_i|^{2+\delta_1}] \leq K$; (iv) $\sigma_d^2 \geq K^{-1}$; (v) $(d_i)_{i \leq n}$ is independent of $(\epsilon_i)_{i \leq n}$.

Assumption 2. (i) $\lambda_{\min}(n^{-1} \sum_{i=1}^n w_i w_i^\top) \geq K^{-1} + o_p(1)$; (ii) $\max_{i \leq n} \mathbb{E}[|w_i|^2] \leq K$; (iii) the first component of the vectors w_i , $1 \leq i \leq n$, is a non-zero constant.

Assumption 3. (i) $n^{-1} \sum_{i=1}^n w_i \epsilon_i = o_p(1)$; (ii) $n^{-1} \sum_{i=1}^n \epsilon_i^2 = n^{-1} \sum_{i=1}^n \mathbb{E}[\epsilon_i^2] + o_p(1)$; (iii) there exists a constant $\delta_2 > 0$ such that $\max_{i \leq n} \mathbb{E}[|\epsilon_i|^{2+\delta_2}] \leq K$; (iv) $n^{-1} \sum_{i=1}^n \mathbb{E}[\epsilon_i^2] \geq K^{-1}$.

Assumption 1 contains our main requirements for the regressor of interest. In particular, Assumption 1(i) holds when $(d_i)_{i \leq n}$ are independently generated from a common distribution. This condition is satisfied in randomized controlled trials (RCTs) that employ i.i.d. assignment. It fails, however, under designs based on individual randomization with fixed treatment arm sizes, as well as under randomized group or cluster assignment, both of which mechanically induce correlation among the d_i s. Assumption 1(ii), which imposes independence between $(d_i)_{i \leq n}$ and $(w_i^\top)_{i \leq n}$, simplifies the development of the main results in this section. This condition holds in many RCTs when $(d_i)_{i \leq n}$ are drawn independently of $(w_i^\top)_{i \leq n}$, but it may fail when treatment assignments are randomized conditional on (a subset of) $(w_i^\top)_{i \leq n}$, as in stratified or covariate-adaptive designs. On the other hand, Assumption 1(ii) can be relaxed; see the remark following Corollary 1 below. Assumptions 1(iii) and 1(iv) are mild moment conditions, where the $2 + \delta_1$ moment requirement in Assumption 1(iii) is mainly used to derive the asymptotic normality of $\hat{\beta}$ via a martingale central limit theorem. Assumption 1(v) means that $(d_i)_{i \leq n}$ is strongly exogenous.

Assumption 2 contains our main requirements for the controls. In particular, Assumption 2(i) means that there is no multicollinearity among controls. Assumption 2(ii) is a mild moment condition. Assumption 2(iii) means that we study regressions with an intercept. Assumption 3 contains our main requirements for the regression error.

Assumption 3(i) holds if ε_i 's are uncorrelated with w_i 's and a law of large numbers applies to the product $w_i \varepsilon_i$. Assumption 3(ii) is essentially a law of large numbers for ε_i^2 's. Assumptions 3(iii) and 3(iv) are mild moment conditions. We highlight that our assumptions allow for a wide array of dependence structures in the matrix $(\varepsilon_i, w_i^\top)_{i \leq n}$, with the main condition in this regard intuitively being that dependence be “weak” enough for the law of large numbers imposed in Assumptions 3(i) and 3(ii) to apply.

For all $1 \leq i \leq n$, denote $d_i^* \equiv d_i - \mu_d$. The following theorem derives the asymptotic distribution of the OLS estimator $\hat{\beta}$ in the strongly exogenous case.

Theorem 1. *Let Assumptions 1, 2 and 3 hold. Then*

$$\frac{\sqrt{n}(\hat{\beta} - \beta)}{\sigma_\varepsilon / \sigma_d} = \frac{n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i}{\sigma_d \sigma_\varepsilon} + o_P(1) \rightarrow_d N(0, 1),$$

where $\sigma_\varepsilon^2 \equiv n^{-1} \sum_{i=1}^n \text{Var}(\varepsilon_i)$.

This theorem establishes two key facts. First, it shows that, given the strong exogeneity of $(d_i)_{i \leq n}$, our mild requirements on the dependence structure of $(\varepsilon_i, w_i^\top)_{i \leq n}$ suffice for establishing asymptotic normality of $\hat{\beta}$. To establish such a conclusion, we rely on a martingale construction that generalizes our discussion in Section 2. Second, Theorem 1 establishes that the asymptotic variance of $\hat{\beta}$ is not affected by the possible correlation across vectors $w_i \varepsilon_i$ since it only depends on the variance of d_i and the averaged variance of the error terms $(\varepsilon_i)_{i \leq n}$. We emphasize that neither of these conclusions need hold for the estimator $\hat{\gamma}$ of the coefficient γ corresponding to the vectors of controls w_i .

In addition, Theorem 1 suggests that we can estimate the variance of $\hat{\beta}$ by $s^2(\check{\mathbf{D}}^\top \check{\mathbf{D}})^{-1}$, where

$$s^2 \equiv n^{-1}(\mathbf{Y} - \mathbf{D}\hat{\beta} - \mathbf{W}\hat{\gamma})^\top (\mathbf{Y} - \mathbf{D}\hat{\beta} - \mathbf{W}\hat{\gamma}), \quad (11)$$

$\check{\mathbf{D}} \equiv \mathbf{M}_W \mathbf{D}$ and $\mathbf{M}_W \equiv \mathbf{I}_n - \mathbf{W}(\mathbf{W}^\top \mathbf{W})^{-1} \mathbf{W}^\top$. The following corollary confirms this conjecture.

Corollary 1. *Let Assumptions 1, 2 and 3 hold. Then*

$$s^2(n^{-1} \check{\mathbf{D}}^\top \check{\mathbf{D}})^{-1} = \sigma_\varepsilon^2 / \sigma_d^2 + o_p(1) \quad \text{and} \quad \frac{\hat{\beta} - \beta}{\sqrt{s^2(\check{\mathbf{D}}^\top \check{\mathbf{D}})^{-1}}} \rightarrow_d N(0, 1). \quad (12)$$

Together with Theorem 1, this corollary is our first main result. Indeed, it is well-known that the top left element of the matrix $(\mathbf{X}^\top \mathbf{X})^{-1}$ coincides with $(\check{\mathbf{D}}^\top \check{\mathbf{D}})^{-1}$. Therefore, this corollary justifies using the classic standard error formula $s^2(\mathbf{X}^\top \mathbf{X})^{-1}$ for inference on β , even under general dependence structures in the errors $(\varepsilon_i)_{i \leq n}$ and controls $(w_i^\top)_{i \leq n}$.

REMARK. Theorem 1 and Corollary 1 can be extended to allow for dependence between d_i and w_i . Indeed, suppose that d_i depends linearly on w_i :

$$d_i = w_i^\top \alpha + \eta_i,$$

where η_i 's satisfy the conditions of Assumption 1 imposed on d_i . By arguments that are similar to those in the proof of Theorem 1 and Corollary 1, it is then straightforward to show that

$$\frac{\sqrt{n}(\hat{\beta} - \beta)}{\sigma_\varepsilon/\sigma_\eta} \rightarrow_d N(0, 1),$$

where σ_η^2 denotes the variance of η_i 's, and that the convergence results (12) still hold. Moreover, σ_ε^2 can be estimated by the same estimator s^2 in (11), and σ_η^2 can be estimated by $n^{-1} \sum_{i=1}^n \hat{\eta}_i^2$ for $\hat{\eta}_i$ residual from regressing $(x_i)_{i \leq n}$ on $(w_i^\top)_{i \leq n}$. $\square$

REMARK. Theorem 1 and Corollary 1 can also be extended to instrumental variable (IV) estimators. Indeed, suppose that

$$d_i = v_i \rho + w_i^\top \alpha + \eta_i, \quad (13)$$

where v_i is an instrumental variable satisfying the conditions of Assumption 1 imposed on d_i and η_i is a (first-stage) estimation error satisfying the conditions of Assumption 3 imposed on ε_i . The two-stage least squares (2SLS) estimator of β then satisfies

$$\hat{\beta}_{2sls} = \frac{\mathbf{V}^\top \mathbf{M}_W \mathbf{Y}}{\mathbf{V}^\top \mathbf{M}_W \mathbf{D}}.$$

where $\mathbf{V} \equiv (v_i)_{i \leq n}$. Using Assumptions 2 and 3, it is then possible to show that

$$\sqrt{n}(\hat{\beta}_{2sls} - \beta) = \frac{n^{-1/2} \sum_{i=1}^n (v_i - \mu_v) \varepsilon_i}{\rho \sigma_v^2} + o_p(1),$$

where μ_v is the mean of v_i 's and σ_v^2 is the variance of v_i 's. Therefore,

$$\frac{\sqrt{n}(\hat{\beta}_{2sls} - \beta)}{\sigma_\varepsilon/(\rho \sigma_v)} \rightarrow_d N(0, 1).$$

It is clear that σ_ε^2 can be estimated by the same s^2 as that in (11) with $\hat{\beta}$ replaced by $\hat{\beta}_{2sls}$, ρ can be estimated by OLS on the (first-stage) regression (13), and σ_v^2 can be estimated by $n^{-1} \tilde{\mathbf{V}}^\top \tilde{\mathbf{V}}$, where $\tilde{\mathbf{V}} = \mathbf{V} - \mathbf{1}_{n,1}(n^{-1} \mathbf{V}^\top \mathbf{1}_{n \times 1})$. $\square$

3.2 Case of Conditional Heteroscedasticity

Next, we derive the asymptotic variance for $\hat{\beta}$ in the case of conditional heteroscedasticity, i.e. when $(\varepsilon_i)_{i \leq n}$ is conditionally heteroscedastic with respect to $(d_i)_{i \leq n}$. Following the notation introduced in Section 2, we focus on the case in which $\varepsilon_i \equiv \sum_{l \leq L} \sigma_l(d_i) \varepsilon_{l,i}^*$, where $e_i \equiv (\varepsilon_{l,i}^*)_{l \leq L}$ is a vector with mean zero, for all $1 \leq i \leq n$.

Let $A_i \equiv (d_i - \mu_d)(\sigma_l(d_i))_{l \leq L}$ for all $1 \leq i \leq n$ and observe that under Assumption 1(i), the random vectors A_i are i.i.d. Denote their common mean vector by μ_A . In addition, denote $\sigma_{e,1}^2 \equiv n^{-1} \sum_{i=1}^n \mathbb{E}[(A_i - \mu_A)^\top e_i]^2$ and $\sigma_{e,2}^2 \equiv \mathbb{E}[(n^{-1/2} \sum_{i=1}^n \mu_A^\top e_i)^2]$. Within this context, we impose the following assumptions.

Assumption 4. (i) $(d_i)_{i \leq n}$ is independent of $(e_i)_{i \leq n}$; (ii) the functions σ_l , $1 \leq l \leq L$, are bounded; (iii) $\sigma_{e,1}^2 \geq K^{-1}$.

Assumption 5. (i) $\|n^{-1} \sum_{i=1}^n e_i e_i^\top - n^{-1} \sum_{i=1}^n \mathbb{E}[e_i e_i^\top]\| = o_p(1)$; (ii) there exists a constant $\delta_3 > 0$ such that $n^{-1} \sum_{i=1}^n \|e_i\|^{2+\delta_3} = O_p(1)$; (iii) $\sigma_{e,2}^{-1} n^{-1/2} \sum_{i=1}^n \mu_A^\top e_i \rightarrow_d N(0, 1)$.

Assumption 4 is mainly used to replace Assumption 1(v) and accounts for the conditional heteroscedasticity of $(\varepsilon_i)_{i \leq n}$ with respect to $(d_i)_{i \leq n}$. Assumption 4(i) requires that $(d_i)_{i \leq n}$ are strongly exogenous with respect to the “scaled” error vector $(e_i)_{i \leq n}$ – a requirement that, as discussed in Section 2 maps well into a potential outcome framework with heterogeneous treatment effects. This condition may fail, for example, when d_i and ε_i exhibit higher-order dependence. Assumption 4(ii) imposes upper bounds on the functions $\sigma_l^2(\cdot)$, which we view as a mild regularity condition. Assumption 4(iii) is a mild moment condition. Assumption 5 contains further restrictions on the vectors e_i . Assumption 5(i) is essentially a law of large numbers for $e_i e_i^\top$ ’s. Assumption 5(ii) is essentially a moment condition for the random variables $\|e_i\|$. Assumption 5(iii) limits the amount of dependence among the vectors e_i to ensure convergence in distribution.

The following theorem, which is our second main result, derives the asymptotic distribution of the OLS estimator $\hat{\beta}$ in the case of conditional heteroscedasticity.

Theorem 2. *Let Assumptions 1(i)-(iv), 2, 3, 4, and 5 hold. Then*

$$\frac{\sqrt{n}(\hat{\beta} - \beta)}{\sigma_{d\varepsilon}/\sigma_d^2} = \frac{n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i}{\sigma_{d\varepsilon}} + o_p(1) \rightarrow_d N(0, 1), \quad (14)$$

where $\sigma_{d\varepsilon}^2 \equiv \sigma_{e,1}^2 + \sigma_{e,2}^2$.

To establish this theorem, we decompose the score $n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i$ into the sum of two uncorrelated terms,

$$n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i = n^{-1/2} \sum_{i=1}^n (A_i - \mu_A)^\top e_i + n^{-1/2} \sum_{i=1}^n \mu_A^\top e_i, \quad (15)$$

and observe that the first term on the right-hand side here forms a martingale difference sequence while the second term is asymptotically normal by assumption. Combining these facts, we are able to obtain asymptotic normality of the sum; see the detailed proof in the Appendix.

Like Theorem 1, this theorem establishes two key facts as well. To see both of them, observe that the term $\sigma_{d\varepsilon}^2$ appearing in the convergence result (14) can be more explicitly rewritten as

$$\sigma_{d\varepsilon}^2 = n^{-1} \sum_{i=1}^n \mathbb{E}[(d_i^*)^2 \varepsilon_i^2] + \mu_A^\top \left[\text{Var} \left(n^{-1/2} \sum_{i=1}^n e_i \right) - n^{-1} \sum_{i=1}^n \text{Var}(e_i) \right] \mu_A. \quad (16)$$

This expression in turn implies that the asymptotic variance of the OLS estimator now depends on the correlation across the vectors e_i , which means that *heteroscedasticity* of the regression

errors forces OLS variance estimators to be adjusted for *correlation* across the errors. On the other hand, the expression (16) also demonstrates that it suffices to adjust the variance estimators only for correlation across the random variables $\mu_A^\top e_i$, instead of correlation across the full vectors e_i . The latter point might seem like a minor technicality but it in fact plays an interesting role in models with heterogeneous treatment effects. Indeed, when $d_i \in \{0, 1\}$ represents the treatment assignment status, $(y_i(1), y_i(0))$ represents the pair of potential outcomes with and without treatment, so that $y_i = y_i(0) + d_i(y_i(1) - y_i(0))$, and w_i consists of a non-zero constant only, it follows that ε_i takes the form (7), which matches the setting here with $e_i = (y_i(0) - \mathbb{E}[y_i(0)], y_i(1) - y_i(0) - \mathbb{E}[y_i(1) - y_i(0)])^\top$ and $\sigma(d_i) = (1, d_i)^\top$. Hence, $\mu_A = (0, \sigma_d^2)^\top$, and so the expression for the asymptotic variance of $\sqrt{n}(\hat{\beta} - \beta)$ reduces to the probability limit of

$$\frac{1}{n\sigma_d^4} \sum_{i=1}^n \mathbb{E}[(d_i - \mu_d)^2 \varepsilon_i^2] + \text{Var} \left(n^{-1/2} \sum_{i=1}^n (y_i(1) - y_i(0)) \right) - n^{-1} \sum_{i=1}^n \text{Var}(y_i(1) - y_i(0)),$$

as previewed in the previous section; see expression (8) there.⁴ In turn, this expression means that it suffices to adjust OLS variance estimation for correlation across treatment effect, and there is no need to worry about correlation of potential outcomes within any given treatment arm. For example, whenever treatment effects $y_i(1) - y_i(0)$ are independent across i , it suffices to use the usual heteroscedasticity-robust variance formulas, even if regression errors ε_i are correlated. Note, however, that it is *necessary* to use heteroscedasticity-robust variance formulas even if the treatment effects are i.i.d. (and not just independent), as the formula $s^2 (\mathbf{X}^\top \mathbf{X})^{-1}$ is not valid in this case. Moreover, the same results apply even if w_i 's include non-constant controls as well, as the second component of e_i 's remains the same in this case.

Finally, we note that if the dependence structure of $\{e_i\}_{i \leq n}$ is known so that the variance of $\sum_{i=1}^n x_i \varepsilon_i$, denoted by V_ε , has an explicit form, then a standard error for $\hat{\beta}$ based on Theorem 2 can, in principle, be constructed as

$$\left(\ell_1^\top (\mathbf{X}^\top \mathbf{X})^{-1} \hat{V}_\varepsilon (\mathbf{X}^\top \mathbf{X})^{-1} \ell_1 \right)^{1/2}, \quad (17)$$

where $\ell_1 \equiv (1, 0_{1 \times d_\gamma})^\top$ and $\hat{V}_\varepsilon$ denotes a consistent estimator of V_ε . However, as is evident from the expression of $\sigma_{d\varepsilon}^2$ in (1), the explicit form of V_ε depends on the specific dependence structure of $\{e_i\}_{i \leq n}$.

As an illustration, suppose that $\{e_i\}_{i \leq n}$ follows a covariance-stationary time series with weak

⁴Equation (8) simplifies to the asymptotic variance formula of the design-based unadjusted average treatment effect estimator derived in Lin (2013) for the binary treatment case, once we adopt a change of notation to reflect the assumption that all randomness in the model arises solely from random assignment. The key simplification is that the second term in (8)—namely, $\text{Var}(n^{-1/2} \sum_{i=1}^n (y_i(1) - y_i(0)))$, which stems from the term $n^{-1/2} \sum_{i=1}^n \mu_A^\top e_i$ in (15), reduces to zero. This is because $n^{-1/2} \sum_{i=1}^n \mu_A^\top e_i$ is non-random under the design-based framework.

dependence, and let $\hat{\varepsilon}_i \equiv y_i - x_i^\top \hat{\theta}$. In this case, V_ε can be estimated using a HAC estimator,

$$\hat{V}_{\varepsilon, HAC} \equiv \sum_{h=-n+1}^{n-1} \mathcal{K}(h/M) \hat{\Gamma}_n(h), \quad (18)$$

where

$$\hat{\Gamma}_n(h) \equiv \sum_{i=h+1}^n x_i x_{i-h}^\top \hat{\varepsilon}_i \hat{\varepsilon}_{i-h} \text{ for } h \geq 0, \text{ and } \hat{\Gamma}_n(h) = \hat{\Gamma}_n(-h)^\top \text{ for } h < 0.$$

Here $\mathcal{K}(\cdot)$ denotes a kernel function and M is a bandwidth. The standard error of $\hat{\beta}$ can then be estimated using (17) with $\hat{V}_\varepsilon$ replaced by $\hat{V}_{\varepsilon, HAC}$. As another example, suppose that $\{e_i\}_{i \leq n}$ exhibits clustered dependence such that e_i and e_j are correlated if and only if i and j belong to the same cluster I_g , where $\{I_g\}_{g \leq G}$ is a collection of mutually exclusive clusters whose union equals $\{1, \dots, n\}$. In this case,

$$\text{Var} \left(\sum_{i=1}^n e_i \right) = \sum_{g \leq G} \text{Var} \left(\sum_{i \in I_g} e_i \right).$$

Accordingly, V_ε can be estimated by

$$\hat{V}_{\varepsilon, CR} \equiv \sum_{g \leq G} \left(\sum_{i \in I_g} x_i \hat{\varepsilon}_i \right) \left(\sum_{i \in I_g} x_i^\top \hat{\varepsilon}_i \right). \quad (19)$$

Therefore, the standard error of $\hat{\beta}$ can be estimated using (17) with $\hat{V}_\varepsilon$ replaced by $\hat{V}_{\varepsilon, CR}$. Consistency of these standard error estimators follows from standard arguments establishing the consistency of HAC estimators (see, e.g., Andrews (1991)) and cluster-robust variance estimators (see, e.g., Hansen (2007)).⁵

4 Group-Level Randomization

A key assumption behind the results of Section 3 is that the regressor of interest is independent and identically distributed across units. In randomized controlled trials, such an assumption is satisfied for completely randomized assignments but can fail under other randomization protocols. For instance, in randomized controlled trials the i.i.d. assumption on the treatment fails when treatment is assigned at a group level. Motivated by this challenge, we suppose in this section that

$$y_{i,j} = d_j \beta + w_{i,j}^\top \gamma + \varepsilon_{i,j}, \quad (20)$$

where the index $j = 1, \dots, n_g$ denotes the group membership, the index $i = 1, \dots, n_j$ denotes units within group j , n_g is the number of groups, n_j is the number of units within group j , d_j denotes

⁵The finite-sample performance of these standard error estimators is examined through simulation studies in Section 5.

the regressor of interest which is invariant within group j , $w_{i,j}$ denotes the vector of controls, and $\varepsilon_{i,j}$ is a mean-zero regression error. We continue to assume that the first component of each $w_{i,j}$ is a non-zero constant and also continue to employ n as the total number of observations, $n = \sum_{j \leq n_g} n_j$.

We emphasize at this point that the index j distinguishes the level at which the regressor d_j varies, but has no special significance for other variables. In particular, we do not insist that any (potential) clustering structure of the errors $((\varepsilon_{i,j})_{i \leq n_j})_{j \leq n_g}$ be the same as the group structure specified by the regressor d_j . In other words, $((\varepsilon_{i,j})_{i \leq n_j})_{j \leq n_g}$ may have a dependence structure completely different from group structure determined by j – e.g., ε_{i_1, j_1} may be correlated with ε_{i_2, j_2} when $j_1 \neq j_2$. This discussion will be formalized in the nature of the regularity conditions below, which do not include, for example, the random effects type specification as in Moulton (1986). Instead, as in the previous section, we will rely on a martingale structure based on $(d_j)_{j \leq n_g}$ that will be crucial for understanding the asymptotic distribution of $\hat{\beta}$.

Following our discussion in the previous section, we consider cases of strong exogeneity and conditional heteroscedasticity separately. We first consider the case of strong exogeneity. In order to study the asymptotic properties of $\hat{\beta}$, we first need to revise Assumptions 1 and 3 to account for the group structure and a possible within-group correlation. To this end, we let $\kappa_n \equiv (\sum_{j \leq n_g} n_j^2)^{1/2}$ and $S_\varepsilon^2 \equiv n^{-1} \sum_{j \leq n_g} \mathbb{E}[(\sum_{i \leq n_j} \varepsilon_{i,j})^2]$ and impose the following assumptions.

Assumption 6. (i) The random variables d_j , $1 \leq j \leq n_g$, are i.i.d. with mean μ_d and variance σ_d^2 ; (ii) $(d_j)_{j \leq n_g}$ is independent of $((w_{i,j}^\top)_{i \leq n_j})_{j \leq n_g}$; (iii) $\max_{j \leq n_g} \mathbb{E}[|d_j|^4] \leq K$; (iv) $\sigma_d^2 \geq K^{-1}$; (v) $(d_j)_{i \leq n_g}$ is independent of $((\varepsilon_{i,j})_{i \leq n_j})_{j \leq n_g}$.

Assumption 7. (i) $n^{-1} \sum_{j=1}^{n_g} \sum_{i=1}^{n_j} w_{i,j} \varepsilon_{i,j} = o_p(S_\varepsilon n^{1/2} / \kappa_n)$; (ii) $S_\varepsilon^{-2} n^{-1} \sum_{j=1}^{n_g} (\sum_{i=1}^{n_j} \varepsilon_{i,j})^2 \rightarrow_p 1$; (iii) there exists a constant $\delta_4 \in (0, 2]$ such that $n^{-1-\delta_4/2} \sum_{j=1}^{n_g} |\sum_{i=1}^{n_j} \varepsilon_{i,j}|^{2+\delta_4} = o_p(S_\varepsilon^{2+\delta_4})$; (iv) $\kappa_n/n = o(1)$.

Assumption 6 requires mild moment restrictions and that the regressor of interest d_j be strongly exogenous in the sense that it be independent of the errors and other regressors. In particular, Assumptions 6(i)–(ii) are satisfied in randomized controlled trials when group assignments are independently drawn from a common distribution. However, these conditions may fail, for example, if $(d_j)_{j \leq n_g}$ are randomized conditional on (a subset of) $((w_{i,j}^\top)_{i \leq n_j})_{j \leq n_g}$, as such designs generally induce dependence between treatment assignment and observed covariates.

To make sense of Assumption 7, assume that each group j has the same size n_j that is independent of n . In this case, κ_n is of order $\sqrt{n}$ and S_ε^2 is typically of order one. In turn, the latter implies that Assumption 7(i) reduces to $n^{-1} \sum_{j \leq n_g} \sum_{i \leq n_j} w_{i,j} \varepsilon_{i,j} = o_p(1)$, which is similar to Assumption 3(i), and Assumption 7(iii) reduces to $n^{-1-\delta_4/2} \sum_{j=1}^{n_g} \sum_{i=1}^{n_j} |\varepsilon_{i,j}|^{2+\delta_4} = o_p(1)$, which is satisfied as long as $\max_{1 \leq j \leq n_g} \max_{1 \leq i \leq n_j} |\varepsilon_{i,j}|^{2+\delta_4} \leq K$. In addition, Assumption 7(iv) reduces to $n^{-1/2} = o(1)$ and is satisfied automatically and Assumption 7(ii) can be regarded as a law of large

numbers. Thus, Assumption 7 in general requires that the size of groups does not increase too fast. Note also that, as previously claimed, this assumption does not require $((w_{i,j}^\top, \varepsilon_{i,j})_{i \leq n_j})_{j \leq n_g}$ to be independent across j .

We are now ready to derive the asymptotic distribution of the OLS estimator $\hat{\beta}$ in the strongly exogenous case with a group-level assignment. In the statement of the result, we impose Assumption 2, which should be interpreted to hold with $\sum_{j=1}^{n_g} \sum_{i=1}^{n_j}$ in place of $\sum_{i=1}^n$ in Assumption 2(i) and $\max_{j \leq n_g} \max_{i \leq n_j}$ in place of $\max_{i \leq n}$ in Assumption 2(ii). Also, we denote $d_j^* \equiv d_j - \mu_d$ for all $1 \leq j \leq n_g$.

Theorem 3. *Let Assumptions 2, 6 and 7 hold. Then*

$$\frac{\sqrt{n}(\hat{\beta} - \beta)}{S_\varepsilon/\sigma_d} = \frac{n^{-1/2} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} \varepsilon_{i,j}}{\sigma_d S_\varepsilon} + o_p(1) \rightarrow_d N(0, 1). \quad (21)$$

The implications of this theorem are following. First, the OLS estimator $\hat{\beta}$ is asymptotically normally distributed under mild moment conditions and under fairly general assumptions on the dependence structure of $((w_{i,j}^\top, \varepsilon_{i,j})_{i \leq n_j})_{j \leq n_g}$. Second, the asymptotic variance of $\hat{\beta}$ is given by $S_\varepsilon^2/\sigma_d^2$. In particular, since $S_\varepsilon^2 = n^{-1} \sum_{j \leq n_g} \text{Var}(\sum_{i \leq n_j} \varepsilon_{i,j})$, when computing standard errors for $\hat{\beta}$ we need only account for possibly within-group j correlation *even if* $((\varepsilon_{i,j})_{i \leq n_j})_{j \leq n_g}$ is dependent across j . Importantly, the group structure is determined solely by the variables $(d_j)_{j \leq n_g}$ and hence is known, considerably simplifying estimation. For instance, in a randomized controlled trial with constant effects, Theorem 3 implies we may, for example, employ Moulton (1986)'s or Liang and Zeger (1986) formulas clustered at the level at which treatment was assigned. We again emphasize, however, that similar conclusions do not apply for $\hat{\gamma}$ whose standard errors and asymptotic normality may depend on the dependence structure of $((w_{i,j}^\top, \varepsilon_{i,j})_{i \leq n_j})_{j \leq n_g}$ across j .

Next, we consider the case of conditional heteroscedasticity. To cover this case, denote $S_{e,1}^2 \equiv n^{-1} \sum_{j=1}^{n_g} \mathbb{E}[(A_j - \mu_A)^\top \sum_{i=1}^{n_j} e_{i,j}]^2$ and $S_{e,2}^2 \equiv \mathbb{E}[(n^{-1/2} \sum_{i=1}^n \mu_A^\top e_i)^2]$. Note that $S_{e,2}$ here actually coincides with $\sigma_{e,2}$ in the previous section. Within this context, we impose the following assumptions.

Assumption 8. (i) $(d_j)_{j \leq n_g}$ is independent of $((e_{i,j}^\top)_{i \leq n_j})_{j \leq n_g}$; (ii) the functions σ_l , $1 \leq l \leq L$, are bounded; (iii) $S_{e,1}/S_\varepsilon \geq K^{-1}$.

Assumption 9. (i) $\|n^{-1} \sum_{j=1}^{n_g} (\sum_{i=1}^{n_j} e_{i,j})(\sum_{i=1}^{n_j} e_{i,j})^\top - n^{-1} \sum_{j=1}^{n_g} \mathbb{E}[(\sum_{i=1}^{n_j} e_{i,j})(\sum_{i=1}^{n_j} e_{i,j})^\top]\| = o_p(S_{e,1}^2)$; (ii) $n^{-2} \sum_{j=1}^{n_g} \|\sum_{i=1}^{n_j} e_{i,j}\|^4 = o_p(S_{e,1}^4)$; (iii) $S_{e,2}^{-1} n^{-1/2} \sum_{i=1}^n \mu_A^\top e_i \rightarrow_d N(0, 1)$.

These assumptions naturally extend Assumptions 4 and 5 in the previous section to allow for group-level assignments. The next theorem derives the asymptotic distribution of the OLS estimator $\hat{\beta}$ in the case of conditional heteroscedasticity with a group-level assignment.

Theorem 4. *Let Assumptions 2, 6(i)-(iv), 7, 8, and 9 hold. Then*

$$\frac{\sqrt{n}(\hat{\beta} - \beta)}{S_{d\varepsilon}/\sigma_d^2} = \frac{n^{-1/2} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} \varepsilon_{i,j}}{S_{d\varepsilon}} + o_p(1) \rightarrow_d N(0, 1), \quad (22)$$

where $S_{d\varepsilon} \equiv S_{e,1}^2 + S_{e,2}^2$.

This theorem relates to Theorem 3 in the same way as Theorem 2 relates to Theorem 1. In particular, noting that the term $S_{d\varepsilon}^2$ appearing in this theorem can be more explicitly rewritten as

$$\begin{aligned} S_{d\varepsilon}^2 &\equiv n^{-1} \sum_{j=1}^{n_g} \mathbb{E} \left[\left(d_j^* \sum_{i=1}^{n_j} \varepsilon_{i,j} \right)^2 \right] \\ &\quad + \mu_A^\top \left[\text{Var} \left(n^{-1/2} \sum_{j=1}^{n_g} \sum_{i=1}^{n_j} e_{i,j} \right) - n^{-1} \sum_{j=1}^{n_g} \text{Var} \left(\sum_{i=1}^{n_j} e_{i,j} \right) \right] \mu_A, \end{aligned}$$

we conclude that because of conditional heteroscedasticity, variance estimators that are clustered at the group level at which the regressor d_j is assigned may not be valid if the errors $e_{i,j}$ are correlated across groups j . On the other hand, in the context of estimation with heterogeneous treatment effects, such estimators are valid if the treatment effects are uncorrelated across these groups.

5 Simulation Studies

In this section, we conduct simulation studies using data from Karlan and List (2007), which examines the effectiveness of matching grants on charitable giving in a large-scale natural field experiment. The experiment was implemented through direct-mail solicitations to 50,083 prior donors of a U.S. nonprofit organization. Individuals were randomly assigned at the individual level to either a control group or a treatment group that received a matching grant offer, with additional randomization over three match ratios, the maximum size of the match, and suggested donation amounts. Embedded in an actual fund raising campaign, the study measured outcomes both as a binary indicator of giving and as the dollar amount donated. The authors find that announcing a matching grant significantly increases both the probability of giving and the average revenue per solicitation.

In our simulations, we abstract from the additional layers of randomization in the original experiment, such as variation in match ratios, maximum match amounts, and suggested contribution levels, and focus exclusively on the primary treatment–control assignment. We further concentrate on the linear regression specification in which the dollar amount donated serves as the dependent variable. This approach isolates the core randomized treatment effect and allows us to use the empirical distribution of outcomes and regression residuals from the original data set to calibrate the

data-generating process, thereby evaluating the finite-sample properties of alternative estimators and inference procedures.⁶

Specifically, we consider the linear regression model specified in (10), where y_i in this simulation study denotes the dollar amount donated by donor i , d_i indicates whether donor i receives a matching offer (with $d_i = 1$ if a matching grant is offered and $d_i = 0$ otherwise), and w_i is a vector that includes a constant term and observed characteristics of donor i . We examine specifications both with and without additional observed characteristics. Accordingly, the regressor vector w_i may consist solely of a constant term or of a constant together with observed donor characteristics.⁷

To simulate the data, we first use the full sample to run a linear regression of y_i on x_i to obtain

$$y_i = x_i^\top \tilde{\theta}_* + \tilde{\varepsilon}_i, \quad (23)$$

where $\tilde{\theta}_* \equiv (\tilde{\beta}_*, \tilde{\gamma}_*^\top)^\top$ denotes the OLS estimator of the unknown coefficient vector $\theta = (\beta, \gamma^\top)^\top$. The regression residuals $\tilde{\varepsilon}_i$ are saved together with the control variables w_i for each donor i , yielding the collection $\{(\tilde{\varepsilon}_i, w_i^\top)\}_{i \leq n}$ which serves as the empirical basis for generating simulated samples.⁸

For each simulation replication s with simulation sample size n_{sim} , we first generate n_{sim} i.i.d. random variables $d_{s,i} \sim \text{Bernoulli}(p)$, where $p = 0.67$ is chosen to match the randomization probability in Karlan and List (2007). The simulated data set $\{(y_{s,i}, d_{s,i}, w_{s,i}^\top)\}_{i \leq n_{\text{sim}}}$ is then constructed using $\{d_{s,i}\}_{i \leq n_{\text{sim}}}$ together with a sampled collection $\{(\tilde{\varepsilon}_{s,i}, w_{s,i}^\top)\}_{i \leq n_{\text{sim}}}$ drawn from $\{(\tilde{\varepsilon}_i, w_i^\top)\}_{i \leq n}$, according to

$$y_{s,i} = x_{s,i}^\top \tilde{\theta}_* + \sigma(d_{s,i}) \tilde{\varepsilon}_{s,i}, \quad \text{for } i = 1, \dots, n_{\text{sim}}, \quad (24)$$

where $\sigma(d_{s,i}) = 1$ corresponds to the conditionally homoscedastic case, while $\sigma(d_{s,i}) = 1 + d_{s,i}$ induces conditional heteroscedasticity,

For both the homoscedastic and heteroscedastic designs, we consider three alternative methods for generating $\{(\tilde{\varepsilon}_{s,i}, w_{s,i}^\top)\}_{i \leq n_{\text{sim}}}$.

1. **Block sampling.** Draw a contiguous block of size n_{sim} from $\{(\tilde{\varepsilon}_i, w_i^\top)\}_{i \leq n}$, thereby preserving the original joint dependence structure.

⁶The data used in the simulation studies are obtained from the Harvard Dataverse repository; see, Karlan and List (2014).

⁷There are 15 observed characteristics included as additional controls in the linear regression, covering donors' past activities (e.g., highest previous contribution and number of prior donations), census demographics (e.g., proportion white and average household size), and several state- and county-level variables. Detailed descriptions of these variables are provided in Table 1 of Karlan and List (2007).

⁸To understand the standard error formula used in Karlan and List (2007), we replicated Table 4, Panel A of that paper; the replication results are reported in Table 3 of Appendix C. We find that they use the homoskedastic standard error presented in (29), which allows us to reproduce all the reported standard errors in Panel A of their Table 4. The only noticeable exception is the standard error of the intercept in column (8), which appears to be a typographical error.

2. **Serial correlation sampling.** Draw $(\tilde{\varepsilon}_{s,i}^*, w_{s,i}^{*\top})$ i.i.d. with replacement from $\{(\tilde{\varepsilon}_i, w_i^\top)\}_{i \leq n}$ for $i = 1, \dots, n_{\text{sim}} + 200$. Generate

$$\tilde{\varepsilon}_{s,i} = 0.5\tilde{\varepsilon}_{s,i-1} + \tilde{\varepsilon}_{s,i}^*, \quad \text{for } i \geq 2 \quad (25)$$

with $\tilde{\varepsilon}_{s,1} = 0$. After discarding the first 200 observations to reduce initialization effects, use $\{(\tilde{\varepsilon}_{s,i}, w_{s,i}^\top)\}_{201 \leq i \leq n_{\text{sim}} + 200}$ in (24), where $w_{s,i} = w_{s,i}^*$.

3. **Cluster dependence sampling.** Draw $(\tilde{\varepsilon}_{s,i}^*, w_{s,i}^{*\top})$ i.i.d. with replacement from $\{(\tilde{\varepsilon}_i, w_i^\top)\}_{i \leq n}$ for $i = 1, \dots, n_{\text{sim}} + 200$, and partition the first n_{sim} observations sequentially into G^* equal-sized clusters $\{I_g^*\}_{g \leq G^*}$, where I_g^* denotes the index set of cluster g . Independently draw cluster shocks $\{\tilde{\varepsilon}_{s,g}^*\}_{g \leq G^*}$ with replacement from $\{\tilde{\varepsilon}_{s,i}^*\}_{n_{\text{sim}}+1 \leq i \leq n_{\text{sim}}+200}$, and set

$$\tilde{\varepsilon}_{s,i} = \tilde{\varepsilon}_{s,i}^* + 0.2\tilde{\varepsilon}_{s,g}^*, \quad \text{for any } i \in I_g^*. \quad (26)$$

The simulated sample is then constructed via (24), with $w_{s,i} = w_{s,i}^*$.

For each simulated sample $\{(y_{s,i}, x_{s,i}^\top)\}_{i \leq n_{\text{sim}}}$, we estimate (10) and obtain

$$y_{s,i} = x_{s,i}^\top \hat{\theta}_s + \hat{\varepsilon}_{s,i}, \quad \text{for } i = 1, \dots, n_{\text{sim}}. \quad (27)$$

where $\hat{\theta}_s$ denotes the OLS estimator and $\hat{\varepsilon}_{s,i}$ the regression residuals. We conduct a two-sided t -test of

$$H_0 : \beta = \tilde{\beta}_*, \quad \text{v.s.} \quad H_1 : \beta \neq \tilde{\beta}_*. \quad (28)$$

We consider three standard error estimators for the coefficient of interest.

The first is the homoscedastic/textbook standard error,

$$\text{SE}_1 \equiv (\hat{\sigma}_{\varepsilon_s}^2 \ell_1^\top (X_s^\top X_s)^{-1} \ell_1)^{1/2}, \quad (29)$$

where $\ell_1 \equiv (1, 0_{1 \times d_\gamma})^\top$, $X_s \equiv (x_{s,1}, \dots, x_{s,n_{\text{sim}}})^\top$ and $\hat{\sigma}_{\varepsilon_s}^2 \equiv (n_{\text{sim}} - d_x)^{-1} \sum_{i \leq n_{\text{sim}}} \hat{\varepsilon}_{s,i}^2$. The second estimator is a HAC standard error obtained from (17) with $\hat{V}_\varepsilon$ replaced by $\hat{V}_{\varepsilon, \text{HAC}}$ in (18). Using the Bartlett kernel, it is given by

$$\text{SE}_2 \equiv \left(\ell_1^\top (X_s^\top X_s)^{-1} \left(\sum_{|h| \leq M} \left(1 - \frac{|h|}{M+1} \right) \hat{\Gamma}_{s,n}(h) \right) (X_s^\top X_s)^{-1} \ell_1 \right)^{1/2}, \quad (30)$$

where

$$\hat{\Gamma}_{s,n}(h) \equiv \sum_{i=h+1}^{n_{\text{sim}}} x_{s,i} x_{s,i-h}^\top \hat{\varepsilon}_{s,i} \hat{\varepsilon}_{s,i-h} \quad \text{for } h \geq 0, \quad \text{and } \hat{\Gamma}_{s,n}(h) = \hat{\Gamma}_{s,n}(-h)^\top \quad \text{for } h < 0.$$

The third estimator is the cluster-robust standard error, obtained from (17) with $\hat{V}_\varepsilon$ replaced by $\hat{V}_{\varepsilon,RC}$ in (19). Specifically,

$$\text{SE}_3(G) \equiv \left(\ell_1^\top (X_s^\top X_s)^{-1} \sum_{g \leq G} \left(\sum_{i \in I_g} x_{s,i} \hat{\varepsilon}_{s,i} \right) \left(\sum_{i \in I_g} x_{s,i}^\top \hat{\varepsilon}_{s,i} \right) (X_s^\top X_s)^{-1} \ell_1 \right)^{1/2}, \quad (31)$$

where the observations are partitioned sequentially into G equal-sized clusters $\{I_g\}_{g \leq G}$.

In the simulation, we set $G^* = n_{\text{sim}}/100$ and consider $n_{\text{sim}} \in \{1000, 2500, 5000\}$ corresponding to approximately 2%, 5%, and 10% of the full sample size in Karlan and List (2007). For the HAC estimator and the cluster-robust standard error, the tuning parameters are specified as

$$M = c \left(\frac{6\rho^2}{(1 - \rho^2)^2} n_{\text{sim}} \right)^{1/3}, \quad \text{for } c \in \{0, 1, 2\}, \quad (32)$$

where $\rho = 0.5$, and

$$G \in \{10, 25, 50\}, \quad (33)$$

respectively.⁹ The constant c in (32) is introduced to assess the sensitivity of inference based on HAC standard errors to the bandwidth choice. When $c = 0$, we have $M = 0$, and the HAC estimator reduces to the heteroscedasticity-consistent (HC) standard error. For the cluster-robust standard error, the true cluster dependence structure under the cluster-dependence sampling scheme is assumed to be unknown. We therefore consider cluster sizes of 10, 25, and 50 to examine the sensitivity of inference to the clustering specification. By construction of the data-generating process, the clusters are correctly specified for

$$(n_{\text{sim}}, G) \in \{(1000, 10), (2500, 25), (5000, 50)\}. \quad (34)$$

We denote by $\text{SE}_2(c)$ the HAC standard error obtained when M is determined according to (32) for a given $c \in \{0, 1, 2\}$. Empirical rejection probabilities of the corresponding t -tests, based on 100,000 Monte Carlo replications, are reported in Tables 1 and 2 for the conditional homoscedastic and heteroscedastic cases, respectively.

From Table 1, it is clear that under conditional homoscedasticity, inference based on the homoscedastic (textbook) standard error, SE_1 , exhibits very good size control across all sampling schemes and sample sizes considered, regardless of whether additional control variables are included in the regression. By contrast, the HAC-based standard error displays mild over-rejection when the sample size is relatively small and/or the bandwidth parameter is relatively large—for example, when $n_{\text{sim}} = 1000$ and/or $c = 2$. The cluster-robust standard error shows nontrivial over-rejection, particularly when the number of clusters is small. Including additional controls

⁹The formula in (32) follows equations (6.2) and (6.4) in Andrews (1991) for the Bartlett kernel, assuming ρ is known.

Table 1: Inference under Conditional Homoskedasticity

	Extra Controls Excluded			Extra Controls Included		
	Block	Serial	Cluster	Block	Serial	Cluster
Panel A: $n = 1000$						
SE ₁	0.0424	0.0461	0.0423	0.0423	0.0465	0.0430
SE ₂ (0)	0.0644	0.0562	0.0629	0.0633	0.0577	0.0619
SE ₂ (1)	0.0682	0.0469	0.0659	0.0673	0.0483	0.0657
SE ₂ (2)	0.0728	0.0519	0.0705	0.0712	0.0529	0.0696
SE ₃ (10)	0.0919	0.0853	0.0905	0.0887	0.0847	0.0872
SE ₃ (25)	0.0743	0.0664	0.0721	0.0714	0.0664	0.0692
SE ₃ (50)	0.0687	0.0605	0.0663	0.0668	0.0608	0.0642
Panel B: $n = 2500$						
SE ₁	0.0462	0.0492	0.0467	0.0471	0.0490	0.0473
SE ₂ (0)	0.0560	0.0533	0.0554	0.0570	0.0537	0.0554
SE ₂ (1)	0.0586	0.0507	0.0577	0.0596	0.0505	0.0573
SE ₂ (2)	0.0604	0.0532	0.0597	0.0611	0.0533	0.0598
SE ₃ (10)	0.0848	0.0825	0.0825	0.0837	0.0827	0.0845
SE ₃ (25)	0.0666	0.0647	0.0663	0.0668	0.0641	0.0660
SE ₃ (50)	0.0619	0.0582	0.0606	0.0610	0.0585	0.0592
Panel C: $n = 5000$						
SE ₁	0.0474	0.0482	0.0485	0.0495	0.0485	0.0482
SE ₂ (0)	0.0530	0.0514	0.0534	0.0548	0.0528	0.0528
SE ₂ (1)	0.0539	0.0498	0.0546	0.0564	0.0510	0.0539
SE ₂ (2)	0.0551	0.0513	0.0562	0.0569	0.0530	0.0550
SE ₃ (10)	0.0830	0.0819	0.0838	0.0848	0.0834	0.0824
SE ₃ (25)	0.0643	0.0621	0.0638	0.0654	0.0638	0.0634
SE ₃ (50)	0.0585	0.0561	0.0590	0.0595	0.0578	0.0572

Note: This table reports the empirical rejection probabilities of two-sided tests of the hypotheses in (28), using the homoskedastic standard error (SE₁), HAC-based standard errors (SE₂(c), with $c = 0, 1, 2$), and cluster-robust standard errors (SE₃(G), with $G = 10, 25, 50$), as defined in (29)–(31). Here SE₂(c) denotes the HAC standard error obtained when M is determined according to (32) for a given $c \in \{0, 1, 2\}$. The labels “Block,” “Serial,” and “Cluster” in the table header correspond to the block sampling, serial correlation, and cluster dependence schemes, respectively. The nominal significance level is 0.05, and critical values are taken from the 0.975 quantile of the standard normal distribution. All results are based on 100,000 Monte Carlo replications.

Table 2: Inference under Conditional Heteroskedasticity

	Extra Controls Excluded			Extra Controls Included		
	Block	Serial	Cluster	Block	Serial	Cluster
Panel A: $n = 1000$						
SE ₁	0.0445	0.0577	0.0749	0.0437	0.0561	0.0732
SE ₂ (0)	0.0529	0.0927	0.0975	0.0536	0.0944	0.0997
SE ₂ (1)	0.0563	0.0906	0.0921	0.0568	0.0915	0.0941
SE ₂ (2)	0.0600	0.0949	0.0890	0.0611	0.0956	0.0909
SE ₃ (10)	0.0813	0.1244	0.0906	0.0794	0.1221	0.0908
SE ₃ (25)	0.0632	0.1037	0.1068	0.0619	0.1036	0.1068
SE ₃ (50)	0.0579	0.0979	0.1017	0.0561	0.0967	0.1017
Panel B: $n = 2500$						
SE ₁	0.0263	0.0400	0.0588	0.0265	0.0403	0.0589
SE ₂ (0)	0.0470	0.0811	0.0975	0.0486	0.0828	0.0999
SE ₂ (1)	0.0491	0.0677	0.0859	0.0508	0.0682	0.0882
SE ₂ (2)	0.0511	0.0695	0.0793	0.0527	0.0698	0.0814
SE ₃ (10)	0.0775	0.1145	0.1273	0.0771	0.1145	0.1289
SE ₃ (25)	0.0569	0.0940	0.0697	0.0582	0.0931	0.0702
SE ₃ (50)	0.0519	0.0869	0.1028	0.0529	0.0878	0.1029
Panel C: $n = 5000$						
SE ₁	0.0207	0.0352	0.0549	0.0207	0.0341	0.0561
SE ₂ (0)	0.0441	0.0786	0.1029	0.0454	0.0791	0.1037
SE ₂ (1)	0.0454	0.0603	0.0876	0.0465	0.0592	0.0896
SE ₂ (2)	0.0466	0.0611	0.0799	0.0476	0.0598	0.0814
SE ₃ (10)	0.0751	0.1106	0.1342	0.0747	0.1100	0.1358
SE ₃ (25)	0.0555	0.0902	0.1142	0.0563	0.0909	0.1145
SE ₃ (50)	0.0497	0.0849	0.0656	0.0499	0.0837	0.0664

Note: This table reports the empirical rejection probabilities of two-sided tests of the hypotheses in (28), using the homoskedastic standard error (SE₁), HAC-based standard errors (SE₂(c), with $c = 0, 1, 2$), and cluster-robust standard errors (SE₃(G), with $G = 10, 25, 50$), as defined in (29)–(31). Here SE₂(c) denotes the HAC standard error obtained when M is determined according to (32) for a given $c \in \{0, 1, 2\}$. The labels “Block,” “Serial,” and “Cluster” in the table header correspond to the block sampling, serial correlation, and cluster dependence schemes, respectively. The nominal significance level is 0.05, and critical values are taken from the 0.975 quantile of the standard normal distribution. All results are based on 100,000 Monte Carlo replications.

does not materially affect the size properties of SE_1 , and it neither meaningfully improves nor worsens the size distortions of the HAC or cluster-robust standard errors. These results are consistent with our theoretical findings: when d_i is randomly assigned so that Assumptions 1(i, ii) hold, inference based on the homoscedastic standard error remains valid even under arbitrary residual dependence.

In the conditional heteroscedastic case, however, the homoscedastic standard error SE_1 loses size control. As shown in Table 2, it tends to under-reject under block and serial-correlation sampling, and to over-reject under cluster dependence. The HAC standard error performs reasonably well in the serial-correlation setting when the sample size is sufficiently large—for example, when $n_{\text{sim}} = 5000$ and c is set to 1, as recommended by Andrews (1991). The cluster-robust standard error generally over-rejects, although its size is closer to the nominal level under correct cluster specification and improves as the sample size increases. These results echo the theoretical conclusions in Subsection 3.2: the homoscedastic standard error is invalid under conditional heteroscedasticity, and valid inference requires explicitly accounting for the dependence structure of the error term. The simulations also provide finite-sample support for the large-sample validity of the variance estimators in (17), which take the forms (30) and (31) under serial correlation and cluster dependence, respectively.

6 Concluding Remarks

This paper shows that when treatment is randomly assigned and independent of other variables, the conventional homoscedastic variance formula delivers valid inference even under arbitrary residual dependence. We characterize the conditions under which this result holds and provide the appropriate variance correction when it does not. We conclude by highlighting a potential application of our findings to shift-share (“Bartik”) designs and by noting a limitation of our main results in nonlinear models.

First, consider the regression

$$y_k^* = \sum_{i=1}^n \omega_{k,i} d_i \beta + u_k, \quad (35)$$

where $\sum_{i=1}^n \omega_{k,i} d_i$ is the shift-share regressor, $\omega_{k,i}$ represents the share of sector i in region k , and d_i denotes a shock or shift in sector i . As emphasized by Adao, Kolesár, and Morales (2019), inference on β is challenging because the residuals u_k may exhibit cross-regional correlation induced by a similar shift-share structure. However, under suitable independence assumptions on the shocks, Adao, Kolesár, and Morales (2019) show that the distribution of the OLS estimator conditional on the residuals does not depend on their correlation structure.

Our results clarify this insight. Writing (35) in matrix form as

$$\mathbf{Y}^* = \Psi \mathbf{D} \beta + \mathbf{U},$$

where $\mathbf{Y}^* \equiv (y_k^*)_{k \leq K}$, $\Psi \equiv (\omega_{k,i})_{k \leq K, i \leq n}$ and $\mathbf{U} \equiv (u_k)_{k \leq K}$, and premultiplying by $(\Psi^\top \Psi)^{-1} \Psi^\top$, we obtain

$$\mathbf{Y} = \mathbf{D}\beta + \mathbf{e} \quad (36)$$

where $\mathbf{Y} \equiv (\Psi^\top \Psi)^{-1} \Psi^\top \mathbf{Y}^*$ and $\mathbf{e} \equiv (\Psi^\top \Psi)^{-1} \Psi^\top \mathbf{U}$. The regressors in (36) are the shocks $\mathbf{D}$, which are assumed independent of the shares and the original residuals $\mathbf{U}$, and hence also of $\mathbf{e}$. By Theorem 1, if the shocks are i.i.d. with zero mean, the OLS estimator in the transformed model is consistent and asymptotically normal (under regularity conditions), regardless of the dependence structure in $\mathbf{U}$. Notably, this estimator differs from the OLS estimator in (35) studied by Adao, Kolesár, and Morales (2019) and Borusyak, Hull, and Jaravel (2022).

A fully rigorous extension would require addressing technical issues such as the triangular-array structure induced by the transformation. Nevertheless, the analysis in Sections 3-4 suggests that extensions to settings with additional controls, instrumental-variable interpretations of the shift-share regressor, or clustered assignment of shocks are conceptually straightforward.

Second, our main results are established for linear models and the OLS estimator of the coefficient on the randomly assigned regressor d_i . In practice, RCT studies may use nonlinear models such as Probit or Logit (see, e.g., Karlan and List (2007) and Falk (2007)). In such models, the martingale structure of the score function that underlies our linear results generally no longer holds. As a result, even when d_i is randomly assigned and independent of other variables, the latent dependence structure of the data may need to be explicitly accounted for in variance estimation.

Overall, our findings provide a simple and transparent benchmark for inference with randomly assigned regressors and point to several promising directions for future research.

References

- ABADIE, A., S. ATHEY, G. W. IMBENS, AND J. M. WOOLDRIDGE (2023): “When should you adjust standard errors for clustering?,” *The Quarterly Journal of Economics*, 138(1), 1–35.
- ADAO, R., M. KOLESÁR, AND E. MORALES (2019): “Shift-share designs: Theory and inference,” *The Quarterly Journal of Economics*, 134(4), 1949–2010.
- ANDREWS, D. W. K. (1991): “Heteroskedasticity and Autocorrelation Consistent Covariance Matrix Estimation,” *Econometrica*, 59(3), 817–858.
- ARELLANO, M. (1987): “Computing robust standard errors for within-groups estimators,” *Oxford bulletin of Economics and Statistics*, 49(4), 431–434.
- BARRIOS, T., R. DIAMOND, G. IMBENS, AND M. KOLESAR (2012): “Clustering, spacial correlations, and randomization inference,” *Journal of American Statistical Association*, 107, 578–591.

- BLOOM, H. S. (2005): *Learning more from social experiments: Evolving analytic approaches*. Russell Sage Foundation.
- BORUSYAK, K., P. HULL, AND X. JARAVEL (2022): “Quasi-experimental shift-share research designs,” *The Review of Economic Studies*, 89(1), 181–213.
- CONLEY, T. G. (1999): “GMM estimation with cross sectional dependence,” *Journal of econometrics*, 92(1), 1–45.
- DASGUPTA, A. (2008): *Asymptotic Theory of Statistics and Probability*. Springer.
- DUFLO, E., R. GLENNERSTER, AND M. KREMER (2007): “Using randomization in development economics research: A toolkit,” *Handbook of development economics*, 4, 3895–3962.
- FALK, A. (2007): “Gift exchange in the field,” *Econometrica*, 75(5), 1501–1511.
- HALL, P., AND C. C. HEYDE (1980): *Martingale Limit Theory and Its Application*. Academic Press, Inc. New York.
- HANSEN, C. B. (2007): “Asymptotic properties of a robust variance matrix estimator for panel data when T is large,” *Journal of Econometrics*, 141(2), 597–620.
- IBRAGIMOV, R., AND U. K. MÜLLER (2010): “t-Statistic based correlation and heterogeneity robust inference,” *Journal of Business & Economic Statistics*, 28(4), 453–468.
- (2016): “Inference with few heterogeneous clusters,” *Review of Economics and Statistics*, 98(1), 83–96.
- KARLAN, D., AND J. A. LIST (2007): “Does price matter in charitable giving? Evidence from a large-scale natural field experiment,” *American Economic Review*, 97(5), 1774–1793.
- KARLAN, D., AND J. A. LIST (2014): “Replication data for: does price matter in charitable giving? evidence from a large-Scale natural field experiment,” Harvard Dataverse, V4, DOI: 10.7910/DVN/27853, <https://doi.org/10.7910/DVN/27853>.
- LIANG, K.-Y., AND S. ZEGER (1986): “Longitudinal Data Analysis Using Generalized Linear Models,” *Biometrika*, 73, 13–22.
- LIN, W. (2013): “Agnostic notes on regression adjustments to experimental data: Reexamining Freedman’s critique,” *The Annals of Applied Statistics*, pp. 295–318.
- MOULTON, B. R. (1986): “Random group effects and the precision of regression estimates,” *Journal of econometrics*, 32(3), 385–397.
- NEWKEY, W. K., AND K. D. WEST (1987): “A Simple, Positive Semi-Definite, Heteroskedasticity and Autocorrelation Consistent Covariance Matrix,” *Econometrica*, 55(3), 703–708.
- WHITE, H. (2014): *Asymptotic theory for econometricians*. Academic press.

Appendix

A Proof of the Main Results

PROOF OF THEOREM 1. Define $\tilde{\mathbf{D}} \equiv \mathbf{D} - \mathbf{1}_{n \times 1}(n^{-1}\mathbf{D}^\top \mathbf{1}_{n \times 1})$ and $\mathbf{M}_W \equiv \mathbf{I}_n - \mathbf{W}(\mathbf{W}^\top \mathbf{W})^{-1}\mathbf{W}^\top$. Then, given that the matrix $\mathbf{W}$ includes a non-zero constant column by Assumption 2(iii), it follows that $\mathbf{1}_{n \times 1}^\top \mathbf{M}_W = 0$. Therefore, by the Frisch-Waugh-Lovell theorem,

$$\hat{\beta} - \beta = (\mathbf{D}^\top \mathbf{M}_W \mathbf{D})^{-1}(\mathbf{D}^\top \mathbf{M}_W \boldsymbol{\epsilon}) = (\tilde{\mathbf{D}}^\top \mathbf{M}_W \tilde{\mathbf{D}})^{-1}(\tilde{\mathbf{D}}^\top \mathbf{M}_W \boldsymbol{\epsilon}). \quad (37)$$

Also, denoting $\bar{d}_n \equiv n^{-1} \sum_{i=1}^n d_i$, we have

$$\mathbb{E}[|\bar{d}_n - \mu_d|^2] = \sigma_d^2/n \leq K/n = o(1) \quad (38)$$

by Assumption 1(i), and so $\bar{d}_n - \mu_d = o_p(1)$ by Markov's inequality. Hence,

$$\bar{d}_n^2 - \mu_d^2 = (\bar{d}_n - \mu_d)(\bar{d}_n + \mu_d) = o_p(1)$$

by Assumption 1(i) again. In addition, by Assumption 1(i) and Khinchin's law of large numbers,

$$n^{-1} \sum_{i=1}^n d_i^2 = \mathbb{E}[d_i^2] + o_p(1), \quad (39)$$

and so

$$n^{-1} \tilde{\mathbf{D}}^\top \tilde{\mathbf{D}} = \mathbb{E}[d_i^2] - \mu_d^2 + o_p(1) = \sigma_d^2 + o_p(1). \quad (40)$$

Further,

$$n^{-1} \tilde{\mathbf{D}}^\top \mathbf{W} = n^{-1} \sum_{i=1}^n d_i^* w_i^\top - \left(n^{-1} \sum_{i=1}^n d_i^* \right) \left(n^{-1} \sum_{i=1}^n w_i^\top \right). \quad (41)$$

By Assumptions 1(i, ii) and 2(ii), we further obtain

$$\mathbb{E} \left[\left\| n^{-1/2} \sum_{i=1}^n d_i^* w_i \right\|^2 \right] = n^{-1} \sum_{i=1}^n \mathbb{E} [||d_i^* w_i||^2] \leq n^{-1} \sum_{i=1}^n \mathbb{E}[d_i^2] \mathbb{E}[||w_i||^2] \leq K.$$

Hence, by Markov's inequality,

$$n^{-1} \sum_{i=1}^n d_i^* w_i = O_p(n^{-1/2}). \quad (42)$$

Similarly, we can use Assumption 1(i) and 2(ii) to show

$$\left(n^{-1} \sum_{i=1}^n d_i^*\right) \left(n^{-1} \sum_{i=1}^n w_i\right) = O_p(n^{-1/2}), \quad (43)$$

which together with (41) and (42) implies that

$$n^{-1} \tilde{\mathbf{D}}^\top \mathbf{W} = O_p(n^{-1/2}). \quad (44)$$

Combining this result with (40) and using Assumptions 2(i), we then have

$$n^{-1} \tilde{\mathbf{D}}^\top \mathbf{M}_W \tilde{\mathbf{D}} = n^{-1} \tilde{\mathbf{D}}^\top \tilde{\mathbf{D}} - n^{-1} \tilde{\mathbf{D}}^\top \mathbf{W} (n^{-1} \mathbf{W}^\top \mathbf{W})^{-1} n^{-1} \mathbf{W}^\top \tilde{\mathbf{D}} = \sigma_d^2 + o_p(1). \quad (45)$$

For the term $\tilde{\mathbf{D}}^\top \mathbf{M}_W \boldsymbol{\epsilon}$ in the numerator of $\hat{\beta} - \beta$ in (37), we have

$$\begin{aligned} n^{-1/2} \tilde{\mathbf{D}}^\top \mathbf{M}_W \boldsymbol{\epsilon} &= n^{-1/2} \tilde{\mathbf{D}}^\top \boldsymbol{\epsilon} - n^{-1/2} \tilde{\mathbf{D}}^\top \mathbf{W} (\mathbf{W}^\top \mathbf{W})^{-1} \mathbf{W}^\top \boldsymbol{\epsilon} \\ &= n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i - \left(n^{-1/2} \sum_{i=1}^n d_i^*\right) \left(n^{-1} \sum_{i=1}^n \varepsilon_i\right) + o_p(1) \\ &= n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i + o_p(1), \end{aligned} \quad (46)$$

where the second equality follows by the definition of $\tilde{\mathbf{D}}$, (44), and Assumptions 2(i) and 3(i), and the third equality follows by Assumptions 1(i) and 3(i) and Markov's inequality. Therefore,

$$\sqrt{n}(\hat{\beta} - \beta) = \frac{n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i}{\sigma_d^2} + o_p(1) \quad (47)$$

by Assumption 1(iv), and so

$$\frac{\sqrt{n}(\hat{\beta} - \beta)}{\sigma_\varepsilon / \sigma_d} = \frac{n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i}{\sigma_d \sigma_\varepsilon} + o_p(1) \quad (48)$$

by Assumptions 1(i) and 3(iv).

We next derive the asymptotic distribution of $n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i / (\sigma_d \sigma_\varepsilon)$. Let $\mathcal{F}_{i,n}$ denote the filtration generated by $(\boldsymbol{\epsilon}^\top, ((d_j)_{j \leq i})^\top)$. Then by Assumptions 1(i, iv) and 3(iii, iv), $n^{-1/2} d_i^* \varepsilon_i / (\sigma_d \sigma_\varepsilon)$ has finite second moment and

$$\mathbb{E} \left[\frac{n^{-1/2} d_i^* \varepsilon_i}{\sigma_d \sigma_\varepsilon} \middle| \mathcal{F}_{i-1,n} \right] = \frac{n^{-1/2} \varepsilon_i}{\sigma_d \sigma_\varepsilon} \mathbb{E} [d_i^* | \mathcal{F}_{i-1,n}] = \frac{n^{-1/2} \varepsilon_i}{\sigma_d \sigma_\varepsilon} \mathbb{E} [d_i^*] = 0 \quad (49)$$

almost surely, which implies that $n^{-1/2} d_i^* \varepsilon_i / (\sigma_d \sigma_\varepsilon)$ is a martingale difference array with respect to $\mathcal{F}_{i,n}$. Next, observe that Assumptions 1(i, v) and 3(ii, iv) yield

$$\sum_{i=1}^n \mathbb{E} \left[\left(\frac{n^{-1/2} d_i^* \varepsilon_i}{\sigma_d \sigma_\varepsilon} \right)^2 \middle| \mathcal{F}_{i-1,n} \right] = \sigma_\varepsilon^{-2} n^{-1} \sum_{i=1}^n \varepsilon_i^2 \rightarrow_p 1. \quad (50)$$

Moreover for any $\eta > 0$, Assumptions 1(iii, iv, v) and 3(iii, iv) allow us to conclude that for $\delta \equiv \min(\delta_1, \delta_2)$,

$$\begin{aligned}
& \sum_{i=1}^n \mathbb{E} \left[\left(\frac{n^{-1/2} d_i^* \varepsilon_i}{\sigma_d \sigma_\varepsilon} \right)^2 \mathbf{1} \left\{ \left| \frac{n^{-1/2} d_i^* \varepsilon_i}{\sigma_d \sigma_\varepsilon} \right| > \eta \right\} \middle| \mathcal{F}_{i-1, n} \right] \\
& \leq \frac{1}{\eta^\delta} \sum_{i=1}^n \mathbb{E} \left[\left| \frac{n^{-1/2} d_i^* \varepsilon_i}{\sigma_d \sigma_\varepsilon} \right|^{2+\delta} \middle| \mathcal{F}_{i-1, n} \right] \\
& = \frac{1}{\eta^\delta (\sigma_d \sigma_\varepsilon)^{2+\delta} n^{1+\delta/2}} \sum_{i=1}^n \mathbb{E} \left[|d_i^*|^{2+\delta} \right] |\varepsilon_i|^{2+\delta} \leq \frac{K}{\eta^\delta n^{\delta/2}} n^{-1} \sum_{i=1}^n |\varepsilon_i|^{2+\delta} = o_p(1). \tag{51}
\end{aligned}$$

In view of (50) and (51), we can invoke the martingale central limit theorem (see, e.g., Corollary 3.1 in Hall and Heyde (1980)) to conclude that

$$\frac{n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i}{\sigma_d \sigma_\varepsilon} \rightarrow_d N(0, 1). \tag{52}$$

The claim of the theorem follows from combining this result with (48). *Q.E.D.*

PROOF OF COROLLARY 1. We first prove that $\lambda_{\min}(n^{-1} \mathbf{X}^\top \mathbf{X}) \geq K^{-1} + o_p(1)$. To do so, observe that

$$n^{-1} \mathbf{D}^\top \mathbf{D} = \mu_d^2 + \sigma_d^2 + o_p(1),$$

by (39). Also, recalling that $d_i^* = d_i - \mu_d$ for all $1 \leq i \leq n$, we have

$$n^{-1} \mathbf{D}^\top \mathbf{W} = n^{-1} \sum_{i=1}^n d_i w_i^\top = \mu_d n^{-1} \sum_{i=1}^n w_i^\top + n^{-1} \sum_{i=1}^n d_i^* w_i^\top = \mu_d n^{-1} \sum_{i=1}^n w_i^\top + o_p(1),$$

where the last equality follows from (42) in the proof of Theorem 1. Hence,

$$\begin{aligned}
n^{-1} \mathbf{X}^\top \mathbf{X} &= n^{-1} (\mathbf{D}, \mathbf{W})^\top (\mathbf{D}, \mathbf{W}) \\
&= n^{-1} \sum_{i=1}^n (\mu_d, w_i^\top)^\top (\mu_d, w_i^\top) + (\sigma_d, \mathbf{0}_{d_w \times 1}^\top)^\top (\sigma_d, \mathbf{0}_{d_w \times 1}^\top) + o_p(1),
\end{aligned}$$

where d_w is the dimension of the vectors w_i . Now, denote

$$R_n \equiv \min \left(\frac{\sqrt{3 \lambda_{\min}(n^{-1} \sum_{i=1}^n w_i w_i^\top)}}{8 |\mu_d|}, \frac{1}{2} \right)$$

and fix any $a_1 \in \mathbb{R}$ and $a_2 \in \mathbb{R}^{d_w}$ such that $a_1^2 + \|a_2\|^2 = 1$. If $|a_1| > R_n$, then

$$(a_1, a_2^\top) \left(n^{-1} \sum_{i=1}^n (\mu_d, w_i^\top)^\top (\mu_d, w_i^\top) + (\sigma_d, \mathbf{0}_{d_w \times 1}^\top)^\top (\sigma_d, \mathbf{0}_{d_w \times 1}^\top) \right) (a_1, a_2^\top)^\top$$

$$\geq a_1^2 \sigma_d^2 \geq R_n^2 \sigma_d^2 \geq K^{-1}$$

by Assumptions 1(i, iv) and 2(i). If, on the other hand, $|a_1| \leq R_n$, then $\|a_2\|^2 = 1 - a_1^2 \geq 1 - 1/4 = 3/4$, and so

$$|a_1| \leq \frac{\|a_2\| \sqrt{\lambda_{\min}(n^{-1} \sum_{i=1}^n w_i w_i^\top)}}{4|\mu_d|}.$$

The latter in turn implies via Jensen's inequality that

$$\begin{aligned} & (a_1, a_2^\top) \left(n^{-1} \sum_{i=1}^n (\mu_d, w_i^\top)^\top (\mu_d, w_i^\top) + (\sigma_d, \mathbf{0}_{d_w \times 1}^\top)^\top (\sigma_d, \mathbf{0}_{d_w \times 1}^\top) \right) (a_1, a_2^\top)^\top \\ & \geq n^{-1} \sum_{i=1}^n (a_1 \mu_d + a_2^\top w_i)^2 \geq n^{-1} \sum_{i=1}^n (a_2^\top w_i)^2 - 2|a_1 \mu_d| n^{-1} \sum_{i=1}^n |a_2^\top w_i| \\ & \geq n^{-1} \sum_{i=1}^n (a_2^\top w_i)^2 - 2|a_1 \mu_d| \sqrt{n^{-1} \sum_{i=1}^n (a_2^\top w_i)^2} \\ & = \sqrt{n^{-1} \sum_{i=1}^n (a_2^\top w_i)^2} \left(\sqrt{n^{-1} \sum_{i=1}^n (a_2^\top w_i)^2} - 2|a_1 \mu_d| \right) \\ & \geq \|a_2\| \sqrt{\lambda_{\min} \left(n^{-1} \sum_{i=1}^n w_i w_i^\top \right)} \left(\|a_2\| \sqrt{\lambda_{\min} \left(n^{-1} \sum_{i=1}^n w_i w_i^\top \right)} - 2|a_1 \mu_d| \right) \\ & \geq 2^{-1} \|a_2\|^2 \lambda_{\min} \left(n^{-1} \sum_{i=1}^n w_i w_i^\top \right) \geq (3/8) \lambda_{\min} \left(n^{-1} \sum_{i=1}^n w_i w_i^\top \right) \geq K^{-1} \end{aligned}$$

by Assumption 2(i). Hence, it follows that

$$\lambda_{\min} \left(n^{-1} \sum_{i=1}^n (\mu_d, w_i^\top)^\top (\mu_d, w_i^\top) + (\sigma_d, \mathbf{0}_{d_w \times 1}^\top)^\top (\sigma_d, \mathbf{0}_{d_w \times 1}^\top) \right) \geq K^{-1},$$

and so

$$\lambda_{\min}(n^{-1} \mathbf{X}^\top \mathbf{X}) \geq K^{-1} + o_p(1), \quad (53)$$

as required.

Next, we prove that s^2 is consistent for σ_ε^2 . To this end, note that by Assumptions 1(i, v) and 3(iii),

$$\mathbb{E} \left[\left(n^{-1} \sum_{i=1}^n d_i^* \varepsilon_i \right)^2 \right] = n^{-2} \sum_{i=1}^n \mathbb{E} [d_i^{*2} \varepsilon_i^2] \leq K n^{-1}. \quad (54)$$

Hence, by Assumptions 2(iii) and 3(i) and Markov's inequality

$$n^{-1} \mathbf{D}^\top \boldsymbol{\epsilon} = n^{-1} \sum_{i=1}^n d_i^* \varepsilon_i + \mathbb{E}[d_i] n^{-1} \sum_i \varepsilon_i = o_p(1),$$

which combined with Assumption 3(i) further implies that

$$n^{-1}\mathbf{X}^\top \boldsymbol{\epsilon} = n^{-1}(\mathbf{D}, \mathbf{W})^\top \boldsymbol{\epsilon} = o_p(1). \quad (55)$$

Combining this bound with (53) gives

$$\hat{\theta} - \theta = (n^{-1}\mathbf{X}^\top \mathbf{X})^{-1}(n^{-1}\mathbf{X}^\top \boldsymbol{\epsilon}) = o_p(1). \quad (56)$$

In addition, $\mathbf{Y} - \mathbf{X}\hat{\theta} = \boldsymbol{\epsilon} - \mathbf{X}(\hat{\theta} - \theta)$. Therefore,

$$\begin{aligned} s^2 &= n^{-1}(\mathbf{Y} - \mathbf{X}\hat{\theta})^\top (\mathbf{Y} - \mathbf{X}\hat{\theta}) = n^{-1}(\boldsymbol{\epsilon} - \mathbf{X}(\hat{\theta} - \theta))^\top (\boldsymbol{\epsilon} - \mathbf{X}(\hat{\theta} - \theta)) \\ &= n^{-1}\boldsymbol{\epsilon}^\top \boldsymbol{\epsilon} - 2n^{-1}(\hat{\theta} - \theta)^\top \mathbf{X}^\top \boldsymbol{\epsilon} + n^{-1}(\hat{\theta} - \theta)^\top \mathbf{X}^\top \mathbf{X}(\hat{\theta} - \theta) \\ &= \sigma_\epsilon^2 + o_p(1), \end{aligned} \quad (57)$$

where the last equality follows from (55) and (56) and Assumptions 1(i), 2(ii) and 3(ii). This finishes the proof of consistency of s^2 .

Now, defining $\tilde{\mathbf{D}} \equiv \mathbf{D} - \mathbf{1}_{n \times 1}(n^{-1}\mathbf{D}^\top \mathbf{1}_{n \times 1})$ to match the proof of Theorem 1 and recalling that $\check{\mathbf{D}} = \mathbf{M}_W \mathbf{D}$, we have

$$\begin{aligned} n^{-1}\check{\mathbf{D}}^\top \check{\mathbf{D}} &= n^{-1}\tilde{\mathbf{D}}^\top \mathbf{M}_W \tilde{\mathbf{D}} = n^{-1}\tilde{\mathbf{D}}^\top \tilde{\mathbf{D}} - n^{-1}(\tilde{\mathbf{D}}^\top \mathbf{W})(\mathbf{W}^\top \mathbf{W})^{-1}(\mathbf{W}^\top \tilde{\mathbf{D}}) \\ &= n^{-1}\tilde{\mathbf{D}}^\top \tilde{\mathbf{D}} + o_p(1) = \sigma_d^2 + o_p(1), \end{aligned}$$

where the third equality follows from (44) and Assumption 2(i), and the fourth from (40). Together with (57), this gives the first convergence result in (12) since $\sigma_d^2 \geq K^{-1}$ by Assumption 1(iv). The second convergence result follows from the first one, Theorem 1, and Slutsky's theorem. *Q.E.D.*

PROOF OF THEOREM 2. As in the proof of Theorem 1, we have

$$\sqrt{n}(\hat{\beta} - \beta) = \frac{n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i}{\sigma_d^2} + o_p(1); \quad (58)$$

see Equation (47) there and note that the derivation of (47) did not rely on Assumption 1(v), which we are not imposing here. Since $\sigma_{d\varepsilon}^2 \geq \sigma_{e,1}^2 \geq K^{-1}$ by Assumption 4(iii) and $\sigma_d^2 \leq K$ by Assumption 1(i), (58) implies that

$$\frac{\sqrt{n}(\hat{\beta} - \beta)}{\sigma_{d\varepsilon}/\sigma_d^2} = \frac{n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i}{\sigma_{d\varepsilon}} + o_p(1),$$

which yields the equality in (14).

We next derive the convergence result in (14), i.e. $\sigma_{d\varepsilon}^{-1} n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i \rightarrow_d N(0, 1)$. To do so, we write

$$n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i = n^{-1/2} \sum_{i=1}^n A_i^\top e_i = n^{-1/2} \sum_{i=1}^n (A_i - \mu_A)^\top e_i + n^{-1/2} \sum_{i=1}^n \mu_A^\top e_i$$

and denote the first and the second terms on the right-hand side by M_1 and M_2 , respectively. Also, we denote $\tilde{A}_i \equiv A_i - \mu_A$ for all $1 \leq i \leq n$. In addition, denote $\hat{\sigma}_{e,1}^2 \equiv n^{-1} \sum_{i=1}^n \mathbb{E}[(\tilde{A}_i^\top e_i)^2 \mid \mathcal{F}_e]$, where $\mathcal{F}_e$ is the filtration generated by $(e_i^\top)_{i \leq n}$. Then

$$\begin{aligned} \hat{\sigma}_{e,1}^2 &= n^{-1} \sum_{i=1}^n e_i^\top \mathbb{E}[\tilde{A}_i \tilde{A}_i^\top] e_i = n^{-1} \sum_{i=1}^n \text{tr}(\mathbb{E}[\tilde{A}_i \tilde{A}_i^\top] e_i e_i^\top) = \text{tr} \left(n^{-1} \sum_{i=1}^n \mathbb{E}[\tilde{A}_i \tilde{A}_i^\top] e_i e_i^\top \right) \\ &= \text{tr} \left(n^{-1} \sum_{i=1}^n \mathbb{E}[\tilde{A}_i \tilde{A}_i^\top] \mathbb{E}[e_i e_i^\top] \right) + o_p(1) \\ &= n^{-1} \sum_{i=1}^n \mathbb{E}[e_i^\top \tilde{A}_i \tilde{A}_i^\top e_i] + o_p(1) = \sigma_{e,1}^2 + o_p(1), \end{aligned} \quad (59)$$

where the first equality follows from Assumption 4(i), the second and the third from properties of the trace operator $\text{tr}(\cdot)$, the fourth from Assumptions 1(i, iii), 4(ii), and 5(i), the fifth from Assumption 4(i) and properties of the trace operator $\text{tr}(\cdot)$, and the sixth from the definition of $\sigma_{e,1}^2$.

Next, let (Z_1, Z_2) be a pair of independent standard normal random variables that is independent of everything else. Then for $\delta = \min(\delta_1, \delta_3)$,

$$\begin{aligned} \sup_{t \in \mathbb{R}} \left| \mathbb{P} \left(\hat{\sigma}_{e,1}^{-1} n^{-1/2} \sum_{i=1}^n \tilde{A}_i^\top e_i \leq t \mid \mathcal{F}_e \right) - \mathbb{P}(Z_1 \leq t) \right| \\ \leq \frac{n^{-1-\delta/2} \sum_{i=1}^n \mathbb{E}[|\tilde{A}_i^\top e_i|^{2+\delta} \mid \mathcal{F}_e]}{\hat{\sigma}_{e,1}^{2+\delta}} \leq \frac{K n^{-1-\delta/2} \sum_{i=1}^n \|e_i\|^{2+\delta}}{\hat{\sigma}_{e,1}^{2+\delta}} = o_p(1), \end{aligned} \quad (60)$$

where the first inequality follows from a version of the Berry-Esseen theorem (see Section 35.1.9 in DasGupta (2008)), the second inequality from the Cauchy-Schwarz inequality and Assumptions 1(iii) and 4(ii), and the last bound from (59) and Assumptions 4(iii) and 5(ii). Therefore, for any $t \in \mathbb{R}$,

$$\begin{aligned} \mathbb{P}(\sigma_{d\varepsilon}^{-1}(M_1 + M_2) \leq t) \\ &= \mathbb{E}[\mathbb{P}(\hat{\sigma}_{e,1}^{-1} M_1 \leq \hat{\sigma}_{e,1}^{-1}(\sigma_{d\varepsilon} t - M_2) \mid \mathcal{F}_e)] = \mathbb{E}[\mathbb{P}(Z_1 \leq \hat{\sigma}_{e,1}^{-1}(\sigma_{d\varepsilon} t - M_2) \mid \mathcal{F}_e)] + o(1) \\ &= \mathbb{P}(\hat{\sigma}_{e,1} Z_1 + M_2 \leq \sigma_{d\varepsilon} t) + o(1) = \mathbb{P}(\sigma_{e,1} Z_1 + M_2 \leq \sigma_{d\varepsilon} t) + o(1) \\ &= \mathbb{E}[\mathbb{P}(\sigma_{e,2}^{-1} M_2 \leq \sigma_{e,2}^{-1}(\sigma_{d\varepsilon} t - \sigma_{e,1} Z_1) \mid Z_1)] + o(1) \\ &= \mathbb{E}[\mathbb{P}(Z_2 \leq \sigma_{e,2}^{-1}(\sigma_{d\varepsilon} t - \sigma_{e,1} Z_1) \mid Z_1)] + o(1) \\ &= \mathbb{P}(\sigma_{d\varepsilon}^{-1}(\sigma_{e,1} Z_1 + \sigma_{e,2} Z_2) \leq t) + o(1) = \mathbb{P}(Z_1 \leq t) + o(1), \end{aligned} \quad (61)$$

where the first equality follows from the law of iterated expectations (LIE), the second from (60) by noting that the difference of two probabilities is always a number between zero and one to conclude that $o_p(1)$ in (60) satisfies $\mathbb{E}[o_p(1)] = o(1)$, the third from the LIE, the fourth from (59) and Assumption 4(iii), the fifth from the LIE, the sixth from Assumption 5(iii), the seventh from

the LIE, and the eighth from noting that $\sigma_{e,1}Z_1 + \sigma_{e,2}Z_2$ is a normal random variable with mean zero and variance $\sigma_{d\varepsilon}^2 = \sigma_{e,1}^2 + \sigma_{e,2}^2$. This gives $\sigma_{d\varepsilon}^{-1}n^{-1/2} \sum_{i=1}^n d_i^* \varepsilon_i \rightarrow_d N(0, 1)$ and completes the proof of the theorem. Q.E.D.

PROOF OF THEOREM 3. For this proof, it will be convenient to denote $\mathbf{W} \equiv ((w_{i,j}^\top)_{i \leq n_j})_{j \leq n_g}$, $\mathbf{Y} \equiv ((y_{i,j})_{i \leq n_j})_{j \leq n_g}$, and $\boldsymbol{\epsilon} \equiv ((\varepsilon_{i,j})_{i \leq n_j})_{j \leq n_g}$. In addition, denote $\mathbf{D} \equiv (d_j \mathbf{1}_{n_j \times 1})_{j \leq n_g}$ and $\tilde{\mathbf{D}} \equiv \mathbf{D} - (n^{-1} \mathbf{D}^\top \mathbf{1}_{n \times 1})$. Moreover, denote $\bar{d}_n \equiv n^{-1} \sum_{j=1}^{n_g} n_j d_j$.

As in the proof of Theorem 1, given that the matrix $\mathbf{W}$ includes a non-zero constant column by Assumption 2(iii), it follows that $\mathbf{1}_{n \times 1}^\top \mathbf{M}_W = 0$. Therefore, by the Frisch-Waugh-Lovell theorem,

$$\hat{\beta} - \beta = (\mathbf{D}^\top \mathbf{M}_W \mathbf{D})^{-1} (\mathbf{D}^\top \mathbf{M}_W \boldsymbol{\epsilon}) = (\tilde{\mathbf{D}}^\top \mathbf{M}_W \tilde{\mathbf{D}})^{-1} (\tilde{\mathbf{D}}^\top \mathbf{M}_W \boldsymbol{\epsilon}). \quad (62)$$

We first consider the denominator $\tilde{\mathbf{D}}^\top \mathbf{M}_W \tilde{\mathbf{D}}$. Since $n = \sum_{j \leq n_g} n_j$, under Assumptions 6(i,iii) we have

$$\mathbb{E} [|\bar{d}_n - \mu_d|^2] = n^{-2} \sum_{j=1}^{n_g} n_j^2 \mathbb{E} [(d_j^*)^2] \leq K n^{-2} \sum_{j=1}^{n_g} n_j^2 = K \kappa_n^2 / n^2. \quad (63)$$

We therefore obtain from Markov's inequality that

$$\bar{d}_n - \mu_d = O_p(\kappa_n/n), \quad (64)$$

and so

$$\bar{d}_n^2 - \mu_d^2 = (\bar{d}_n - \mu_d)(\bar{d}_n + \mu_d) = O_p(\kappa_n/n). \quad (65)$$

by Assumption 6(iii). In addition, again under Assumptions 6(i, iii) we have

$$\begin{aligned} \mathbb{E} \left[\left| n^{-1} \sum_{j=1}^{n_g} n_j (d_j^2 - \mathbb{E}[d_j^2]) \right|^2 \right] &= n^{-2} \sum_{j=1}^{n_g} n_j^2 \mathbb{E} [|d_j^2 - \mathbb{E}[d_j^2]|^2] \\ &\leq K n^{-2} \sum_{j=1}^{n_g} n_j^2 = K \kappa_n^2 / n^2, \end{aligned}$$

and so

$$n^{-1} \sum_{j=1}^{n_g} n_j (d_j^2 - \mathbb{E}[d_j^2]) = O_p(\kappa_n/n) \quad (66)$$

by Markov's inequality. Combining results (65) and (66) then yields

$$n^{-1} \tilde{\mathbf{D}}^\top \tilde{\mathbf{D}} = n^{-1} \sum_{j=1}^{n_g} n_j (d_j - \bar{d}_n)^2 = n^{-1} \sum_{j=1}^{n_g} n_j d_j^2 - (\bar{d}_n)^2 = \sigma_d^2 + O_p(\kappa_n/n). \quad (67)$$

Further, since d_j does not depend on i , we have

$$n^{-1} \tilde{\mathbf{D}}^\top \mathbf{W} = n^{-1} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} w_{i,j}^\top - \left(n^{-1} \sum_{j=1}^{n_g} n_j d_j^* \right) \left(n^{-1} \sum_{j=1}^{n_g} \sum_{i=1}^{n_j} w_{i,j}^\top \right)$$

$$= n^{-1} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} w_{i,j}^\top + O_p(\kappa_n/n), \quad (68)$$

where the second equality is by Assumption 2(ii) and (64). Moreover, by Assumptions 2(ii) and 6(i, ii, iii),

$$\mathbb{E} \left[\left\| n^{-1} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} w_{i,j}^\top \right\|^2 \right] = \sigma_d^2 n^{-2} \sum_{j=1}^{n_g} \mathbb{E} \left[\left\| \sum_{i=1}^{n_j} w_{i,j} \right\|^2 \right] \leq K n^{-2} \sum_{j=1}^{n_g} n_j^2 = K \kappa_n^2 / n^2.$$

Therefore, by Markov's inequality we obtain

$$n^{-1} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} w_{i,j}^\top = O_p(\kappa_n/n),$$

which together with (68) further shows that

$$n^{-1} \tilde{\mathbf{D}}^\top \mathbf{W} = O_p(\kappa_n/n). \quad (69)$$

Collecting the results (67), and (69) and using Assumptions 2(i) and 7(iv), we get

$$n^{-1} \tilde{\mathbf{D}}^\top \mathbf{M}_W \tilde{\mathbf{D}} = n^{-1} \tilde{\mathbf{D}}^\top \tilde{\mathbf{D}} - n^{-1} \tilde{\mathbf{D}}^\top \mathbf{W} (n^{-1} \mathbf{W}^\top \mathbf{W})^{-1} n^{-1} \mathbf{W}^\top \tilde{\mathbf{D}} = \sigma_d^2 + o_p(1). \quad (70)$$

Next, we consider the numerator $\tilde{\mathbf{D}}^\top \mathbf{M}_W \boldsymbol{\epsilon}$ in (62). By (64) and Assumptions 7(i) and 2(iii),

$$\frac{\bar{d}_n - \mu_d}{S_\varepsilon} n^{-1/2} \sum_{j=1}^{n_g} \sum_{i=1}^{n_j} \varepsilon_{i,j} = \frac{O_p(\kappa_n/n)}{S_\varepsilon} n^{-1/2} \sum_{j=1}^{n_g} \sum_{i=1}^{n_j} \varepsilon_{i,j} = o_p(1).$$

Therefore,

$$\begin{aligned} \frac{n^{-1/2} \tilde{\mathbf{D}}^\top \boldsymbol{\epsilon}}{S_\varepsilon} &= \frac{n^{-1/2} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} \varepsilon_{i,j}}{S_\varepsilon} - \frac{(\bar{d}_n - \mu_d)}{S_\varepsilon} n^{-1/2} \sum_{j=1}^{n_g} \sum_{i=1}^{n_j} \varepsilon_{i,j} \\ &= \frac{n^{-1/2} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} \varepsilon_{i,j}}{S_\varepsilon} + o_p(1). \end{aligned} \quad (71)$$

In addition,

$$\frac{n^{-3/2} \kappa_n \mathbf{W}^\top \boldsymbol{\epsilon}}{S_\varepsilon} = o_p(1)$$

by Assumption 7(i), which together with (69) and (71) and Assumption 2(i) shows

$$\begin{aligned} \frac{n^{-1/2} \tilde{\mathbf{D}}^\top \mathbf{M}_W \boldsymbol{\epsilon}}{S_\varepsilon} &= \frac{n^{-1/2} \tilde{\mathbf{D}}^\top \boldsymbol{\epsilon}}{S_\varepsilon} - \frac{n^{-1/2} \tilde{\mathbf{D}}^\top \mathbf{W} (\mathbf{W}^\top \mathbf{W})^{-1} \mathbf{W}^\top \boldsymbol{\epsilon}}{S_\varepsilon} \\ &= \frac{n^{-1/2} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} \varepsilon_{i,j}}{S_\varepsilon} + o_p(1). \end{aligned} \quad (72)$$

Given that $S_\varepsilon^{-1}n^{-1/2}\sum_{j=1}^{n_g}(d_j - \mu_d)\sum_{i=1}^{n_j}\varepsilon_{i,j} = O_p(1)$ by Assumptions 6(i, iii) and 7(ii) and Markov's inequality and that $\sigma_d^2 \geq K^{-1}$ by Assumption 6(iv), we obtain from (62), (70), and (72) that

$$\frac{\sqrt{n}(\hat{\beta} - \beta)}{S_\varepsilon} = \frac{n^{-1/2}\sum_{j=1}^{n_g}d_j^*\sum_{i=1}^{n_j}\varepsilon_{i,j}}{\sigma_d^2 S_\varepsilon} + o_p(1)$$

and so

$$\frac{\sqrt{n}(\hat{\beta} - \beta)}{S_\varepsilon/\sigma_d} = \frac{n^{-1/2}\sum_{j=1}^{n_g}d_j^*\sum_{i=1}^{n_j}\varepsilon_{i,j}}{\sigma_d S_\varepsilon} + o_p(1) \quad (73)$$

by Assumption 6(iii).

We next derive the asymptotic distribution of $n^{-1/2}\sum_{j=1}^{n_g}d_j^*\sum_{i=1}^{n_j}\varepsilon_{i,j}/(\sigma_d S_\varepsilon)$. Let $\mathcal{F}_{j,n}$ denote the filtration generated by $(\boldsymbol{\epsilon}^\top, ((d_m)_{m \leq j})^\top)$. Then by Assumptions 6(i, v),

$$\begin{aligned} \mathbb{E} \left[\frac{n^{-1/2}d_j^*\sum_{i=1}^{n_j}\varepsilon_{i,j}}{\sigma_d S_\varepsilon} \middle| \mathcal{F}_{j-1,n} \right] &= \frac{n^{-1/2}\sum_{i=1}^{n_j}\varepsilon_{i,j}}{\sigma_d S_\varepsilon} \mathbb{E} [d_j^* \middle| \mathcal{F}_{j-1,n}] \\ &= \frac{n^{-1/2}\sum_{i=1}^{n_j}\varepsilon_{i,j}}{\sigma_d S_\varepsilon} \mathbb{E} [d_j^*] = 0 \end{aligned}$$

almost surely, which implies that $n^{-1/2}d_j^*\sum_{i=1}^{n_j}\varepsilon_{i,j}/(\sigma_d S_\varepsilon)$ is a martingale difference array with respect to $\mathcal{F}_{j,n}$. Next, observe that Assumptions 6(i, v) and 7(ii) yield

$$\sum_{j=1}^{n_g} \mathbb{E} \left[\left(\frac{n^{-1/2}d_j^*\sum_{i=1}^{n_j}\varepsilon_{i,j}}{\sigma_d S_\varepsilon} \right)^2 \middle| \mathcal{F}_{j-1,n} \right] = S_\varepsilon^{-2}n^{-1}\sum_{j=1}^{n_g} \left(\sum_{i=1}^{n_j}\varepsilon_{i,j} \right)^2 \rightarrow_p 1. \quad (74)$$

Moreover for any $\eta > 0$, Assumptions 6(iii, iv, v) and 7(iii) allow us to conclude that

$$\begin{aligned} &\sum_{j=1}^{n_g} \mathbb{E} \left[\left(\frac{n^{-1/2}d_j^*\sum_{i=1}^{n_j}\varepsilon_{i,j}}{\sigma_d S_\varepsilon} \right)^2 \mathbf{1} \left\{ \left| \frac{n^{-1/2}d_j^*\sum_{i=1}^{n_j}\varepsilon_{i,j}}{\sigma_d S_\varepsilon} \right| > \eta \right\} \middle| \mathcal{F}_{j-1,n} \right] \\ &\leq \frac{1}{\eta^{\delta_4}} \sum_{j=1}^{n_g} \mathbb{E} \left[\left| \frac{n^{-1/2}d_j^*\sum_{i=1}^{n_j}\varepsilon_{i,j}}{\sigma_d S_\varepsilon} \right|^{2+\delta_4} \middle| \mathcal{F}_{j-1,n} \right] \\ &= \frac{1}{\eta^{\delta_4}(\sigma_d S_\varepsilon)^{2+\delta_4}n^{1+\delta_4/2}} \sum_{j=1}^{n_g} \mathbb{E} [|d_i^*|^{2+\delta_4}] \left| \sum_{i=1}^{n_j}\varepsilon_{i,j} \right|^{2+\delta_4} \\ &\leq \frac{K}{\eta^{\delta_4}S_\varepsilon^{2+\delta_4}n^{1+\delta_4/2}} \sum_{j=1}^{n_g} \left| \sum_{i=1}^{n_j}\varepsilon_i \right|^{2+\delta_4} = o_p(1). \end{aligned}$$

Combining this bound with (74), we can invoke the martingale central limit theorem (see, e.g., Corollary 3.1 in Hall and Heyde (1980)) to conclude that

$$\frac{n^{-1/2}\sum_{j=1}^{n_g}d_j^*\sum_{i=1}^{n_j}\varepsilon_{i,j}}{\sigma_d S_\varepsilon} \rightarrow_d N(0, 1). \quad (75)$$

The claim of the theorem follows from combining this result with (73).

Q.E.D.

PROOF OF THEOREM 4. As in the proof of Theorem 3, we have

$$\frac{\sqrt{n}(\hat{\beta} - \beta)}{S_\varepsilon/\sigma_d} = \frac{n^{-1/2} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} \varepsilon_{i,j}}{\sigma_d S_\varepsilon} + o_p(1); \quad (76)$$

see Equation (73) there and note that the derivation of (73) did not rely on Assumption 6(v), which we are not imposing here. Since $S_{d\varepsilon} \geq S_{e,1}$ and $S_{e,1} \geq S_\varepsilon K^{-1}$ by Assumption 8(iii) and $\sigma_d \leq K$ by Assumption 6(iii), (76) implies that

$$\frac{\sqrt{n}(\hat{\beta} - \beta)}{S_{d\varepsilon}/\sigma_d^2} = \frac{n^{-1/2} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} \varepsilon_{i,j}}{S_{d\varepsilon}} + o_p(1),$$

which yields the equality in (22).

We next derive the convergence result in (22), i.e. $S_{d\varepsilon}^{-1} n^{-1/2} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} \varepsilon_{i,j} \rightarrow_d N(0, 1)$. To do so, we write

$$\begin{aligned} n^{-1/2} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} \varepsilon_{i,j} &= n^{-1/2} \sum_{j=1}^{n_g} A_j^\top \sum_{i=1}^{n_j} e_{i,j} \\ &= \sum_{j=1}^{n_g} (A_j - \mu_A)^\top \sum_{i=1}^{n_j} e_{i,j} + n^{-1/2} \sum_{j=1}^{n_g} \mu_A^\top \sum_{i=1}^{n_j} e_{i,j} \end{aligned}$$

and denote the first and the second terms on the right-hand side by M_1 and M_2 , respectively. Also, we denote $\tilde{A}_j \equiv A_j - \mu_A$ for all $1 \leq j \leq n_g$. In addition, denote $\hat{\Sigma}_{e,1}^2 \equiv n^{-1} \sum_{i=1}^n \mathbb{E}[(\tilde{A}_j^\top e_{i,j})^2 \mid \mathcal{F}_e]$, where $\mathcal{F}_e$ is the filtration generated by $((e_{i,j}^\top)_{i \leq n_j})_{j \leq n_g}$.

Then $\hat{S}_{e,1}^2 - S_{e,1}^2 = o_p(S_{e,1}^2)$ by the same argument as that used to derive (59) in the proof of Theorem 2 with Assumption 9(i) replacing Assumption 5. In addition, letting Z be a standard normal random variable that is independent of everything else, we have

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P} \left(\hat{\Sigma}_{e,1}^{-1} n^{-1/2} \sum_{j=1}^{n_g} \tilde{A}_j^\top \sum_{i=1}^{n_j} e_{i,j} \leq t \mid \mathcal{F}_e \right) - \mathbb{P}(Z \leq t) \right| = o_p(1)$$

by the same argument as that used to derive (60) in the proof of Theorem 2 with $\delta = 2$ and Assumption 9(ii) replacing Assumption 5(ii). Finally,

$$\mathbb{P}(S_{d\varepsilon}^{-1} (M_1 + M_2) \leq t) = \mathbb{P}(Z \leq t) + o(1)$$

for all $t \in \mathbb{R}$ by the same argument as that used to derive (61) in the proof of Theorem 2 with Assumption 9(iii) replacing Assumption 5(iii). This gives the convergence result

$$S_{d\varepsilon}^{-1} n^{-1/2} \sum_{j=1}^{n_g} d_j^* \sum_{i=1}^{n_j} \varepsilon_{i,j} \rightarrow_d N(0, 1)$$

and completes the proof of the theorem.

Q.E.D.

B Asymptotic Equivalence of Two Variance Formulae

Lemma 1. *Consider the linear regression model in (1). Suppose: (i) $(d_i)_{i \leq n}$ are i.i.d. with zero mean, finite and nonzero variance σ_d^2 ; (ii) $(\varepsilon_i)_{i \leq n}$ is covariance stationary with auto-covariance function $\Gamma_\varepsilon(\cdot)$ satisfying $\Gamma_\varepsilon(0) > 0$ and $\sum_{j=1}^{\infty} \Gamma_\varepsilon(j)^2 < \infty$. For $\mathbf{D} \equiv (d_i)_{i \leq n}$, we then have*

$$\frac{\mathbf{D}^\top \Omega \mathbf{D}}{\Gamma_\varepsilon(0) \mathbf{D}^\top \mathbf{D}} \rightarrow 1 \text{ almost surely as } n \rightarrow \infty, \quad (77)$$

where Ω is the covariance matrix of $(\varepsilon_i)_{i \leq n}$.

PROOF OF LEMMA 1. Note that

$$n^{-1} \mathbf{D}^\top \Omega \mathbf{D} = n^{-1} \sum_{i_1=1}^n \sum_{i_2=1}^n d_{i_1} d_{i_2} \Gamma_\varepsilon(i_1 - i_2) = \Gamma_\varepsilon(0) n^{-1} \mathbf{D}^\top \mathbf{D} + 2n^{-1} \sum_{i=2}^n U_i \quad (78)$$

where $U_i \equiv \sum_{i'=1}^{i-1} x_i x_{i'} \Gamma_\varepsilon(i - i')$. Let $\tilde{\mathcal{F}}_i$ denote the natural filtration generated by $(d_j)_{j \leq i}$. Then, under the assumption that $(d_i)_{i \leq n}$ is i.i.d., it follows that $\{U_i, \tilde{\mathcal{F}}_i\}$ is a martingale difference sequence with variance $\mathbb{E}[U_i^2] = \sigma_d^4 \sum_{j=1}^{i-1} \Gamma_\varepsilon(j)^2$. Therefore we obtain that

$$\begin{aligned} \sum_{i=2}^n i^{-2} \mathbb{E}[U_i^2] &= \sigma_d^4 \sum_{i=2}^n i^{-2} \sum_{j=1}^{i-1} \Gamma_\varepsilon(j)^2 \\ &= \sigma_d^4 \sum_{j=1}^{n-1} \Gamma_\varepsilon(j)^2 \sum_{m=j+1}^n m^{-2} \leq K \sigma_d^4 \sum_{j=1}^{\infty} \Gamma_\varepsilon(j)^2 < \infty \end{aligned} \quad (79)$$

where the first inequality follows from $\sum_{m=1}^{\infty} m^{-2} < K$ and the last inequality is due to $\sigma_d^2 < \infty$ and $\sum_{j=1}^{\infty} \Gamma_\varepsilon(j)^2 < \infty$ by assumption. Hence, (79) establishes that $\sum_{i=2}^{\infty} i^{-2} \mathbb{E}[U_i^2] < \infty$. By the martingale strong law of large numbers (see, e.g., Theorem 3.76 in White (2014)) we can therefore deduce that $n^{-1} \sum_{i=2}^n U_i \rightarrow 0$ almost surely as $n \rightarrow \infty$, which together with (78) yields

$$n^{-1} \mathbf{D}^\top \Omega \mathbf{D} - \Gamma_\varepsilon(0) n^{-1} \mathbf{D}^\top \mathbf{D} \rightarrow 0 \text{ almost surely as } n \rightarrow \infty. \quad (80)$$

Moreover, by condition (i) of the lemma and Kolmogorov's strong law of large numbers (see, e.g., Theorem 3.1 in White (2014)), $n^{-1} \mathbf{D}^\top \mathbf{D} \rightarrow \sigma_d^2$ almost surely as $n \rightarrow \infty$. Since $\sigma_d^2 > 0$ and $\Gamma_\varepsilon(0) > 0$, the claim of the lemma then follows from (80). Q.E.D.

C Replication of Karlan and List (2007)

Table 3: Replication of Table 4, Panel A, in Karlan and List (2007)

Dependent variable:	Amount given						Change in amount given	
	All		Already gave in 2005		Did not give in 2005		All	
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
<i>d</i>	0.154 (0.083)	0.118 (0.151)	0.152 (0.093)	0.102 (0.170)	0.157 (0.140)	0.142 (0.255)	6.331 (12.014)	-18.650 (21.925)
<i>d</i> * ratio2		0.089 (0.117)		-0.027 (0.132)		0.216 (0.198)		17.945 (16.985)
<i>d</i> * ratio3		0.001 (0.117)		-0.114 (0.132)		0.121 (0.197)		18.583 (16.987)
<i>d</i> * 25,000 maximum		0.045 (0.135)		0.024 (0.152)		0.068 (0.228)		25.256 (19.612)
<i>d</i> * 50,000 maximum		-0.126 (0.135)		0.066 (0.152)		-0.337 (0.228)		25.191 (19.615)
<i>d</i> * 100,000 maximum		-0.111 (0.135)		0.046 (0.152)		-0.289 (0.228)		24.637 (19.612)
<i>d</i> * medium gift		0.090 (0.117)		0.220 (0.132)		-0.053 (0.197)		0.315 (16.985)
<i>d</i> * high gift		0.069 (0.117)		-0.032 (0.132)		0.179 (0.197)		-18.214 (16.987)
Constant	0.813 (0.067)	0.813 (0.067)	0.423 (0.076)	0.423 (0.076)	1.241 (0.114)	1.241 (0.114)	-56.893 (9.810)	-56.893 (9.810)
<i>R</i> -squared	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Observations	50,083	50,083	26,217	26,217	23,866	23,866	50,083	50,083

Note: OLS coefficients are reported. Parentheses contain homoskedastic standard errors $\sqrt{\text{diag}\{\hat{\sigma}_{\varepsilon_s}^2(X_s^\top X_s)^{-1}\}}$.